

REMARKS ON HILBERT IDENTITIES, ISOMETRIC EMBEDDINGS, AND INVARIANT CUBATURE

HIROSHI NOZAKI AND MASANORI SAWA

ABSTRACT. Victor (2004) developed a method to construct cubature formulae with various combinatorial objects. Motivated by this, we generalize Victor's method with one more combinatorial object, called regular t -wise balanced designs. Many cubature of small indices with few points are provided, which are used to update Shatalov's table (2001) of isometric embeddings in small-dimensional Banach spaces, as well as to improve some classical Hilbert identities. A famous theorem of Bajnok (2007) on Euclidean designs invariant under the Weyl group of Lie type B is extended to all finite irreducible reflection groups. A short proof of the Bajnok theorem is presented in terms of Hilbert identities.

1. INTRODUCTION

Let p be a positive integer such that $p \neq \infty$. The m -dimensional Euclidean space $\mathbb{R}^m$ is a Banach space l_p^m endowed with the norm

$$\|x\|_p = \left(\sum_{i=1}^m |x_i|^p \right)^{1/p}.$$

Given two spaces l_p^m and l_q^n , a classical problem in Banach space theory asks when there is an $\mathbb{R}$ -linear map $F: l_p^m \rightarrow l_q^n$ such that

$$\|F(x)\|_q = \|x\|_p$$

for every $x \in l_p^m$. Such a map is called an *isometric embedding from l_p^m to l_q^n* . To exclude trivial cases, we assume that $n \geq m \geq 2$ and $p \neq q$. It is known [22, Theorem 1.1] that if $p, q \neq \infty$ and an isometric embedding from l_p^m to l_q^n exists, then $p = 2$ and q is an even integer. Throughout this paper we only consider the case where $p = 2$ and q is even, and fix the notations p, q, m, n .

Isometric embeddings are closely related to a representation of $(\sum_{i=1}^m x_i^2)^{q/2}$ as a sum of q th powers of linear forms with positive real coefficients. Such representations originally stem from a work of Hilbert on Waring's problem [16], and therefore called *Hilbert identities* [25]. Hilbert solved Waring's problem, showing on the way that there exist isometric embeddings $l_2^m \rightarrow l_q^n$ with n depending on m and q . Several alternative proofs of Hilbert's theorem are known; for example, see [6], [7] and the references therein. But most of them, including the original by Hilbert, involve

1991 *Mathematics Subject Classification.* Primary 65D32, 11E76, Secondary 52A21.

Key words and phrases. Cubature formula, Hilbert identity, isometric embedding, Victor method.

The second author is supported in part by Grant-in-Aid for Young Scientists (B) 22740062 and Grant-in-Aid for Challenging Exploratory Research 23654031 by the Japan Society for the Promotion of Science.

non-constructive arguments in analysis, and do not give any explicit constructions of embeddings¹. Thus publications with explicit embeddings continued to appear.

Isometric embeddings are also related to a certain object in numerical analysis. Let Ω be a subset of $\mathbb{R}^m$ on which a normalized measure μ is defined. A finite subset X of Ω with a positive weight w is called a *cubature formula of index q* if

$$(1.1) \quad \int_{\Omega} f(x) \mu(dx) = \sum_{x \in X} w(x) f(x)$$

for every $f \in \text{Hom}_q(\Omega)$, where $\text{Hom}_q(\Omega)$ is the space of all homogeneous polynomials of degree q restricted to Ω . Lyubich and Vaserstein [22] and Reznick [27] proved the equivalence between an embedding $l_2^m \rightarrow l_q^n$ and an n -point cubature of index q for the surface measure ρ on the $(m-1)$ -dimensional unit sphere S^{m-1} .

Many papers are devoted to the construction of spherical cubature formulae. There are two classical approaches: One uses orbits of finite subgroups of the orthogonal group $O(m)$ acting on S^{m-1} [33], and the other takes “product” of several lower-dimensional cubature [34]. Cubature formulae that are studied in the context of numerical analysis and related areas, are often of degree type. Victoir [35] developed a novel technique to construct degree-type cubature for integrals with special symmetry. His idea is as follows: Given a cubature formula invariant under the Weyl group of Lie type B , one eliminates some specified points of the formula by using combinatorial objects such as t -designs and orthogonal arrays. With this method, Victoir found many cubature of small degrees with few points in general dimensional spaces.

This paper has several important aims. First, we generalize the Victoir method with a special class of block designs, called *regular t -wise balanced designs*. The concept of regular t -wise balanced designs has been substantiated by applications in statistics [8, 10, 18], however, it seems that there is insufficient evidence to support it from other mathematical aspects. To find a new meaning of this concept, as well as to let it know researchers in many areas of mathematics are both important aims of this paper. On the other hand, Bajnok [1, Theorem 3] proved that Euclidean designs, a generalization of spherical cubature, that are invariant under the Weyl group of Lie type B have degree at most 7. We further discuss the Bajnok theorem both from a combinatorial and analytic point of view.

This paper is organized as follows. In Section 2 we review some basic facts and notions, and explain the Victoir method in detail. In Section 3 we generalize the Victoir method with regular t -wise balanced designs. In Section 4, we give general-dimensional index-four and -six cubature, together with some extra examples of index-six cubature that improve Shatalov’s table [32, Theorem 4.7.20] of isometric embeddings $l_2^m \rightarrow l_6^n$. In Section 5, we generalize the Bajnok theorem for all finite irreducible reflection groups, and thereby classify spherical cubature with a certain geometric meaning. In Section 6, some of the cubature constructed in Sections 4 and 5 are translated into Hilbert identities, in order to improve classical identities as such by Schur [6] and Reznick [27]. An extremely short proof of the Bajnok theorem is given in terms of Hilbert identities.

¹ Bruce Reznick kindly told us that Stridsberg’s proof (1912) is constructive, if we know how to compute the roots of Hermite polynomials.

2. PRELIMINARIES

2.1. Isometric embeddings and Hilbert identities. Lyubich and Vaserstein [22] and Reznick [27] observed a close relationship between Hilbert identities, isometric embeddings, and spherical cubature formulae.

Theorem 2.1. *The following are equivalent.*

- (i) *There exists a cubature formula of index q on S^{m-1} with n points;*
- (ii) *There exists an isometric embedding $l_2^m \rightarrow l_q^n$;*
- (iii) *There exist n vectors $r_1, \dots, r_n \in \mathbb{R}^m$ such that for any $x \in \mathbb{R}^m$*

$$\langle x, x \rangle^{\frac{q}{2}} = \sum_{i=1}^n \langle x, r_i \rangle^q.$$

We explain Theorem 2.1 in detail for further arguments in the following sections. Assume that points $x_1, \dots, x_n \in S^{m-1}$ and weights $w_1, \dots, w_n$ form a cubature of index q on S^{m-1} . Let $\langle x, y \rangle^q \in \text{Hom}_q(\mathbb{R}^m)$, where $\langle \cdot, \cdot \rangle$ denotes the usual inner product. Then

$$\sum_{i=1}^n w_i \langle x, x_i \rangle^q = \int_{S^{m-1}} \langle x, y \rangle^q \rho(dy) = \langle x, x \rangle^{\frac{q}{2}} c_q,$$

where

$$c_q = \int_{S^{m-1}} y_1^q \rho(dy), \quad y = (y_1, \dots, y_m).$$

This is, equivalently,

$$\langle x, x \rangle^{\frac{q}{2}} = \sum_{i=1}^n \langle x, r_i \rangle^q,$$

where $r_i = \sqrt[q]{w_i/c_q} x_i$. This polynomial identity is further transformed as follows:

$$\langle x, x \rangle^{\frac{1}{2}} = \left(\sum_{i=1}^n \langle x, r_i \rangle^q \right)^{\frac{1}{q}},$$

which implies that the mapping

$$x \mapsto (\langle x, r_1 \rangle, \dots, \langle x, r_n \rangle)$$

is an isometric embedding $l_2^m \rightarrow l_q^n$.

By the early fundamental works of Hilbert [16], there is a positive integer $N(m, q)$ such that for any $n \geq N(m, q)$ an isometric embedding $l_2^m \rightarrow l_q^n$ exists. It is known (cf. [27]) that

$$(2.1) \quad \binom{m + \frac{q}{2} - 1}{m - 1} \leq N(m, q) \leq \binom{m + q - 1}{m - 1}.$$

The lower- and upper-bound part of (2.1) mean the dimension of $\text{Hom}_{q/2}(\mathbb{R}^m)$ and $\text{Hom}_q(\mathbb{R}^m)$ respectively.

2.2. Cubature formulae. Let $\Omega \subset \mathbb{R}^m$, and μ be a normalized measure on Ω such that Ω, μ are both invariant under the group $O(m)$. We assume that polynomials are integrable up to sufficiently large degrees for

$$\mathcal{I}[f] = \int_{\Omega} f(x) \mu(dx).$$

Let X be a finite set in $\mathbb{R}^m$ with a positive weight w . The pair (X, w) is called a *cubature formula of degree q for $\mathcal{I}$* if

$$\mathcal{I}[f] = \sum_{x \in X} w(x)f(x)$$

for every $f \in \mathcal{P}_q(\Omega)$, where $\mathcal{P}_q(\Omega)$ denotes the space of all polynomials of degree at most q restricted to Ω . In particular, a spherical cubature is called a *spherical design* if w is a constant weight.

A subset X of $\mathbb{R}^m$ is said to be *antipodal* if it is partitioned into $\tilde{X}, -\tilde{X}$, namely, $X = \tilde{X} \cup (-\tilde{X})$ and $\tilde{X} \cap (-\tilde{X}) = \emptyset$. A cubature formula (X, w) is *centrally symmetric* if X is antipodal and $w(x) = w(-x)$ for any $x \in X$. The following mentions the relationship among degree-type and index-type spherical cubature.

Proposition 2.2. ([22, Proposition 4.3]). *Let X be an antipodal finite subset of S^{m-1} . Then X is a centrally symmetric cubature formula on S^{m-1} of degree $q+1$ with $2n$ points iff $\tilde{X}$ is a cubature formula on S^{m-1} of index q with n points.*

We are interested in the following type of integrals:

$$(2.2) \quad \int_{\mathbb{R}^m} f(x)W(\|x\|_2)dx,$$

where W is a density function on $\mathbb{R}^m$. Such integrals are often considered in the context of analysis; for example see [37].

Proposition 2.3. *If points $x_1, \dots, x_n$ and weights $w_1, \dots, w_n$ form a cubature formula of index q for (2.2), then the points $x_i/\|x_i\|_2$ and the weights $\frac{\|x_i\|_2^q w_i}{\int_0^\infty r^{q+m-1}W(r)dr}$ form a cubature formula of index q on S^{m-1} . Conversely, if $x_1, \dots, x_n$ and $w_1, \dots, w_n$ form a cubature formula of index q on S^{m-1} , then the points x_i and weights $w_i \int_0^\infty r^{q+m-1}W(r)dr$ form a cubature formula of index q for (2.2).*

Proof. The result follows by observing that for any $f \in \text{Hom}_q(\mathbb{R}^m)$,

$$\begin{aligned} \int_{\mathbb{R}^m} f(x)W(\|x\|_2)dx &= \int_0^\infty \left(\int_{S^{m-1}} f(rx)\rho(dx) \right) r^{m-1}W(r)dr \\ &= \int_0^\infty r^{q+m-1}W(r)dr \int_{S^{m-1}} f(x)\rho(dx). \end{aligned}$$

□

Remark 2.4. By Proposition 2.3, in order to construct spherical cubature, we may find cubature for any integral of the form (2.2). For example, one may think of Gaussian integrals. Such cubature formulae are of particular interest in probability theory [21] and algebraic combinatorics [2]. Moreover, the m -dimensional Gaussian integral can be represented simply as the m -fold product of one-dimensional Gaussian integrals, which is convenient for explaining Victor's method.

The following proposition is often used in Sections 3 and 4.

Proposition 2.5. *Let X be an antipodal finite subset of R^m . Let $\tilde{w}, w$ be weight functions on $\tilde{X}, X$, respectively, such that for any $x \in \tilde{X}$, $w(x) = w(-x) = 2\tilde{w}(x)$. Then $(\tilde{X}, \tilde{w})$ is a cubature formula of index q for (2.2) if and only if (X, w) is a centrally symmetric cubature of index q for (2.2).*

2.3. The Sobolev theorem. Let G be a finite subgroup of $O(m)$, and $f \in \mathcal{P}_t(\mathbb{R}^m)$. We define the action of $\sigma \in G$ on f as follows:

$$(\sigma f)(x) = f(x^{\sigma^{-1}}), \quad x \in \mathbb{R}^m.$$

A polynomial f is said to be G -invariant if $\sigma f = f$ for every $\sigma \in G$. We denote the set of G -invariant polynomials in $\mathcal{P}_t(\mathbb{R}^m)$, $\text{Harm}_t(\mathbb{R}^m)$ by $\mathcal{P}_t(\mathbb{R}^m)^G$, $\text{Harm}_t(\mathbb{R}^m)^G$ respectively, where $\text{Harm}_t(\mathbb{R}^m)$ is the subspace of $\mathcal{P}_t(\mathbb{R}^m)$ of harmonic homogeneous polynomials of degree t .

A cubature formula is said to be G -invariant if the domain and measure of the integral are invariant under G , the points are a union of G -orbits $z_1^G, \dots, z_e^G$, and $w(x) = w(x')$ for any $x, x' \in z_i^G$; the orbits $z_1^G, \dots, z_e^G$ and weights $w_1, \dots, w_e$ are said to *generate* the formula.

Theorem 2.6 ([33]). *With the above set up, a G -invariant cubature formula is of degree t if and only if it is exact for every polynomial $f \in \mathcal{P}_t(\mathbb{R}^m)^G$.*

Theorem 2.6 is known as the Sobolev theorem, which is at the core of the Victor method, as seen in the next subsection.

The concept of Euclidean designs was introduced by Neumaier and Seidel [23] as a generalization of spherical cubature. Let X be a finite set in $\mathbb{R}^m$, and $\{\|x\|_2 \mid x \in X\} = \{r_1, \dots, r_p\}$. Let S_i^{m-1} be the sphere of radius r_i centered at the origin, and $X_i = X \cap S_i^{m-1}$. To each S_i the surface measure ρ_i is assigned. Let $|S_i^{m-1}| = \int_{S_i^{m-1}} \rho_i(dx)$, where $\frac{1}{|S_i^{m-1}|} \int_{S_i^{m-1}} f(x) \rho_i(dx) = f(0)$ if $S_i^{m-1} = \{0\}$.

Definition 2.7 ([23]). With the above set up, X is a *Euclidean t -design* of $\mathbb{R}^m$ if

$$(2.3) \quad \sum_{i=1}^p \frac{\sum_{x \in X_i} w(x)}{|S_i^{m-1}|} \int_{S_i^{m-1}} f(x) \rho_i(dx) = \sum_{x \in X} w(x) f(x)$$

for every polynomial $f \in \mathcal{P}_t(S)$.

As readily seen by the definition, Euclidean designs can be viewed as cubature formulae on multiple concentric spheres.

The following is a variation of the Sobolev theorem for Euclidean designs, which generalizes the familiar theorem of Neumaier and Seidel [23].

Theorem 2.8 ([24]). *Let G be a subgroup of $O(m)$. Let $X = \cup_{k=1}^M r_k x_k^G$, where $x_k \in S^{m-1}$ and $r_k > 0$. Then the following are equivalent:*

- (i) X is a G -invariant Euclidean t -design of $\mathbb{R}^m$.
- (ii) $\sum_{x \in X} w(x) \|x\|^{2j} \varphi(x) = 0$ for any $\varphi \in \text{Harm}_l(\mathbb{R}^m)^G$, $1 \leq l \leq t$, $0 \leq j \leq \lfloor \frac{t-l}{2} \rfloor$.

Hereafter let G be an irreducible reflection group in $\mathbb{R}^m$. Such groups are completely classified [4]. Let integers $1 = d_1 \leq d_2 \leq \dots \leq d_m$ be the exponents of G (see [4, Ch.V, §6]).

Theorem 2.9 ([9]). *Let G be a finite irreducible reflection group. Let $q_i = \dim(\text{Harm}_i(\mathbb{R}^m)^G)$. Then*

$$\sum_{i=0}^{\infty} q_i \lambda^i = \prod_{i=2}^m \frac{1}{1 - \lambda^{1+d_i}}.$$

In particular, for any $x \in \mathbb{R}^m$, the orbit x^G is a spherical d_2 -design in S^{m-1} .

Let $\alpha_1, \dots, \alpha_m$ be the fundamental roots of a reflection group G . The *corner vectors* $v_1, \dots, v_m$ are defined by $v_i \perp \alpha_j$ if and only if $i \neq j$. We may assume that $\|v_k\|_2 = 1$. We consider the set

$$\mathcal{X}(G, J) = \bigcup_{k \in J} r_k v_k^G,$$

where $J \subset \{1, 2, \dots, m\}$ and $r_k > 0$. Let R denote the set of r_k .

Theorem 2.10. (Bajnok [1, Theorem 3]). *Let $m \geq 2$ be an integer. Then there is no choice of R, J , and w for which $(\mathcal{X}(B_m, J), w)$ is a Euclidean 8-design of $\mathbb{R}^m$.*

Similar results are known for the groups A_{m-1}, D_m [24]. In Section 5, we generalize these results, and determine the maximum degree of invariant Euclidean designs for all irreducible reflection groups.

2.4. The Victor method.

2.4.1. Combinatorial tools. Let K be a set of positive integers $k_1, \dots, k_\ell$. A pair of v elements V , and subsets $\mathcal{B}$ of V of cardinalities from K is called a *t -wise balanced design*, denoted by t -(v, K, λ), if every t elements of V occur exactly λ times in $\mathcal{B}$. Elements of V and $\mathcal{B}$ are called *points* and *blocks*. In particular if K is a singleton, say $K = \{k\}$, a t -wise balanced design is called a *t -design*, and is denoted by t -(v, k, λ). In this paper we only consider designs without repeated blocks.

It is well known (cf. [19]) that for $0 \leq t' \leq t$ and a subset $T' \subset V$ of t' elements, the number of blocks of a t -(v, k, λ) design containing T' is given as

$$(2.4) \quad \lambda \frac{\binom{v-t'}{t-t'}}{\binom{k-t'}{t-t'}} = \frac{(v-t')(v-t'-1) \cdots (v-t+1)}{(k-t')(k-t'-1) \cdots (k-t+1)},$$

not depending on the choice of T' . For each $0 \leq t' \leq t$, a t -design is also a t' -design. In general t -wise balanced designs do not necessarily have this property; see Section 3 for the detail.

Let $(V, \mathcal{B})$ be a t -wise balanced design with v points and b blocks. An incidence matrix M of the design $(V, \mathcal{B})$ is a $v \times b$ zero-one matrix which has a row for each point and a column for each block, and for $x \in V$ and $B \in \mathcal{B}$, (x, B) -entry takes 1 iff $x \in B$. Given real numbers α, β , let $\mathbf{v}_l(\alpha, \beta)$ be a v -dimensional vector such that the first l coordinates are α and the remaining $v-l$ coordinates are β . For example, $\mathbf{v}_l(\alpha, 0)^{B_v}$ means the vertices of a generalized hyperoctahedron that is inscribed in the $(v-1)$ -dimensional sphere of radius $\sqrt{l\alpha^2}$ [1]. To the matrix M , we associate a *generalized incidence matrix with parameters α, β* by defining $I_{\alpha, \beta} = \beta J_{v, b} + (\alpha - \beta)M$, where $\alpha \neq \beta$ and $J_{v, b}$ is the all-one matrix of size $v \times b$.

An $N \times l$ matrix with entries ± 1 is called an *orthogonal array with strength t , constraints l and index λ* , if in every t columns, each of the 2^t ordered combinations of elements ± 1 appears in exactly λ rows. We denote this by $OA(N, l, 2, t)$. We do not put λ in the notation, since $\lambda = N/2^t$ by the definition. When $l \leq t$, we allow *trivial OA*, namely, the $2^l \times l$ matrix such that every 2^l ordered combinations of elements ± 1 appears in exactly one row.

2.4.2. Victor's method. The group B_m contains two special subgroups: the subgroup L of all transpositions of coordinates in $\mathbb{R}^m$, the subgroup $\hat{L}$ of all sign changes which is isomorphic to the elementary abelian 2-group $(\mathbb{Z}/2\mathbb{Z})^m$. It turns out that $|y^{\hat{L}}| = 2^{|\text{wt}(y)|}$, where $\text{wt}(y)$ is the number of nonzero coordinates of a vector y .

We denote by $\mathcal{I}$ the Gaussian integral

$$\mathcal{I}[f] = \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} f((x_1^2, \dots, x_m^2)) \exp(-\frac{\|x\|_2^2}{2}) dx_1 \cdots dx_m.$$

This is equivalent to the integral $\hat{\mathcal{I}}$ on the first orthant $\mathbb{R}_+^m$

$$\hat{\mathcal{I}}[f] = \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}_+^m} f((x_1, \dots, x_m)) \exp(-\frac{\|x\|_1}{2}) (\prod_{i=1}^m x_i)^{-1/2} dx_1 \cdots dx_m.$$

Let

$$x^2 = (x_1^2, \dots, x_m^2)$$

for $x = (x_1, \dots, x_m) \in \mathbb{R}^m$, and

$$\sqrt{x} = (\sqrt{x_1}, \dots, \sqrt{x_m})$$

for $x = (x_1, \dots, x_m) \in \mathbb{R}_+^m$.

Proposition 2.11. (cf. [35, 38]). *If $z_1^{\hat{L}}, \dots, z_e^{\hat{L}}$ and $w_1, \dots, w_e$ generate an $\hat{L}$ -invariant cubature formula of degree q for $\mathcal{I}$, then $z_1^2, \dots, z_e^2$ and $w_1 2^{\text{wt}(z_1)}, \dots, w_e 2^{\text{wt}(z_e)}$ form a cubature formula of degree $q/2$ for $\hat{\mathcal{I}}$. Conversely, if $z_1, \dots, z_e$ and $w_1, \dots, w_e$ form a cubature of degree $q/2$ for $\hat{\mathcal{I}}$, then $\sqrt{z_1^{\hat{L}}}, \dots, \sqrt{z_e^{\hat{L}}}$ and $w_1/2^{\text{wt}(z_1)}, \dots, w_e/2^{\text{wt}(z_e)}$ generate a cubature of degree q for $\mathcal{I}$.*

The following theorem is due to Victor [35, Subsection 4.4].

Theorem 2.12. (i) *Assume that there exist a cubature formula of degree $q/2$ for $\hat{\mathcal{I}}$ of the form*

$$\hat{\mathcal{I}}[f] = \frac{w}{\binom{m}{k}} \sum_{x \in \mathbf{v}_k(\alpha, \beta)^L} f(x) + \sum_{i=1}^M \frac{w_i}{|x_i^L|} \sum_{x \in x_i^L} f(x),$$

and a $q/2$ -design with m points and b blocks of size k . Let X be the columns of a generalized incidence matrix with parameters α, β . Then,

$$\hat{\mathcal{I}}[f] = \frac{w}{b} \sum_{x \in X} f(x) + \sum_{i=1}^M \frac{w_i}{|x_i^L|} \sum_{x \in x_i^L} f(x)$$

is a cubature formula of degree $q/2$.

(ii) *Assume that there exist an $\hat{L}$ -invariant cubature formula of degree q for $\mathcal{I}$ of the form*

$$\mathcal{I}[f] = \sum_{i=1}^M \frac{\lambda_i}{2^{\text{wt}(x_i)}} \sum_{x \in x_i^{\hat{L}}} f(x),$$

and $\text{OA}(|X_i|, \text{wt}(x_i), 2, q)$ with rows X_i for $i = 1, \dots, M$. Then,

$$\mathcal{I}[f] = \sum_{i=1}^M \frac{\lambda_i}{|X_i|} \sum_{x \in X_i} f(x)$$

is a cubature formula of degree q .

The Victoir method was originally written in a more general setting. For example, the integrals considered there are not restricted to Gaussian integrals. In this paper, however, we took only Gaussian integrals since Victoir's ideas can be fully understood with Gaussian integrals.

3. GENERALIZING THE VICTOIR METHOD

In this section we generalize the Victoir method with a strengthening of the concept of t -wise balanced designs. We use the notations $B_m, L, \tilde{L}, \mathcal{I}, \tilde{\mathcal{I}}, \mathbf{v}_i(\cdot, \cdot), \text{wt}(\cdot)$ that are defined in Subsection 2.4.

A t -wise balanced design $(V, \mathcal{B})$ is said to be *regular* if for each $0 \leq t' \leq t$ and each t' -subset T' of V , the number of blocks containing T' does not depend on the choice of T' [8]. As noted in Subsection 2.4, any t -design possesses this property, but t -wise balanced designs do not always so. When $t = 2$, this concept is equivalent to that of *equireplicate* 2-wise balanced designs [10].

Let $\mathcal{B}$ be the set of blocks of a regular t -(v, K, λ) design, where $K = \{k_1, \dots, k_f\}$. Let $\mathcal{B}_i = \{B \in \mathcal{B} \mid |B| = k_i\}$. Let $y_i \in \mathbb{R}^m$ with $\text{wt}(y_i) = k_i$, and $y_K = \{y_1, \dots, y_f\}$. We define the following discrete measure:

$$\delta_{y_K, L} := \sum_{i=1}^f \frac{|\mathcal{B}_i|}{|\mathcal{B}| \binom{m}{k_i}} \sum_{x \in y_i^L} \delta_x.$$

Proposition 3.1. *Assume that there exists a regular t -($m, \{k_i \mid 1 \leq i \leq f\}, \lambda$) design $(V, \mathcal{B})$. Let X be the columns of a generalized incidence matrix with parameters α, β with $\alpha \neq \beta$. Let $y_1, \dots, y_f \in X$ such that $\text{wt}(y_i) = k_i$. Then*

$$\int_{\bigcup_{i=1}^f y_i^L} f(x) \delta_{y_K, L}(dx) = \frac{1}{|\mathcal{B}|} \sum_{x \in X} f(x)$$

for every $f \in \mathcal{P}_t(\bigcup_{i=1}^f y_i^L)$.

Proof. By changing variables $x_i \rightarrow (x_i - \beta)/(\alpha - \beta)$, there is no loss of generality in assuming $\alpha = 1, \beta = 0$. Then for any $e_1, \dots, e_m \geq 0$,

$$\int_{\bigcup_{i=1}^f y_i^L} f(x_1^{e_1}, \dots, x_m^{e_m}) \delta_{y_K, L}(dx) = \int_{\bigcup_{i=1}^f y_i^L} f(x_1, \dots, x_m) \delta_{y_K, L}(dx).$$

Permuting the rows of an incidence matrix also gives another t -wise balanced design with the same parameters $m, k_1, \dots, k_f, \lambda$. Thus it suffices to show that

$$\int_{\bigcup_{i=1}^f y_i^L} f(x) \delta_{y_K, L}(dx) = \frac{1}{|\mathcal{B}|} \sum_{x \in X} f(x)$$

for the monomials $f(x) = \prod_{i=1}^j x_i, 1 \leq j \leq t$. To do this, we count the pairs $(T', B) \in \binom{V}{t'} \times \mathcal{B}$, $T' \subset B$ in two ways:

$$\lambda' \binom{m}{t'} = \sum_{T' \in \binom{V}{t'}} \sum_{T' \subset B \in \mathcal{B}} 1 = \sum_{B \in \mathcal{B}} \sum_{\substack{T' \subset B \\ T' \in \binom{V}{t'}}} 1 = \sum_{i=1}^f \sum_{B \in \mathcal{B}_i} \sum_{\substack{T' \subset B \\ T' \in \binom{V}{t'}}} 1 = \sum_{i=1}^f |\mathcal{B}_i| \binom{k_i}{t'},$$

where the regularity is used to show the first equality. Thus, for $f(x) = \prod_{i=1}^{t'} x_i$,

$$\sum_{y \in X} f(y) = \lambda' = \sum_{i=1}^f \frac{|\mathcal{B}_i| \binom{k_i}{t'}}{\binom{m}{t'}}.$$

This is further transformed to

$$\begin{aligned} \sum_{i=1}^f \frac{|\mathcal{B}_i|}{\binom{m}{k_i}} \cdot \frac{\binom{m}{k_i} \binom{k_i}{t'}}{\binom{m}{t'}} &= \sum_{i=1}^f \frac{|\mathcal{B}_i|}{\binom{m}{k_i}} \cdot \binom{m-t'}{k_i-t'} \\ &= \sum_{i=1}^f \frac{|\mathcal{B}_i|}{\binom{m}{k_i}} \sum_{x \in y_i^L} f(x) = |\mathcal{B}| \cdot \int_{\bigcup_{i=1}^f y_i^L} f(x) \delta_{y_K, L}(\mathrm{d}x). \end{aligned}$$

□

Remark 3.2. In a combinatorial framework (cf. [31]), some researchers regard t -wise balanced designs as cubature on “discrete spheres”. However, among them, there are only a few publications where the regularity of designs is mentioned. Victorio seems to be the first who employed combinatorial t -designs to reduce the size of cubature for ordinary continuous integrals.

The following generalizes Theorem 2.12 (i) and motivates the study of regular t -wise balanced designs both in a combinatorial and analytic manner.

Theorem 3.3. *Assume that there exists a regular $q/2$ -wise balanced design with m points and b_i blocks of size k_i , $i = 1, \dots, e$. Moreover assume that there exists a cubature formula of degree $q/2$ (or index $q/2$) for $\hat{\mathcal{I}}$ of the form*

$$\hat{\mathcal{I}}[f] = c \left(\sum_{i=1}^e \frac{b_i}{\binom{m}{k_i} b} \sum_{x \in \mathbf{v}_{k_i}(\alpha, \beta)^L} f(x) \right) + \sum_{i=2}^M \frac{w_i}{|x_i^L|} \sum_{x \in x_i^L} f(x)$$

where b is the total number of blocks of the design and c is a positive number. Let X be the columns of a generalized incidence matrix with parameters α, β . Then

$$\hat{\mathcal{I}}[f] = \frac{c}{b} \sum_{x \in X} f(x) + \sum_{i=2}^M \frac{w_i}{|x_i^L|} \sum_{x \in x_i^L} f(x)$$

is a cubature formula of degree $q/2$ (or index $q/2$).

The following proposition is often used in Section 4.

Proposition 3.4. *Assume there exists a t -(v, k, λ) design. Then the following hold:*

- (i) *There exists a regular t -($v-1, \{k, k-1\}, \lambda$) design with $\lambda \frac{\binom{v-1}{t-1}}{\binom{k-1}{t-1}}$ blocks of size $k-1$ and $\frac{(v-k)\lambda}{k} \frac{\binom{v-1}{t-1}}{\binom{k-1}{t-1}}$ blocks of size k .*
- (ii) *Let X be the columns of an incidence matrix of the design given in (i), and $y_1 = \mathbf{v}_k(1, 0)$, $y_2 = \mathbf{v}_{k-1}(1, 0)$. Then for every $f \in \mathcal{P}_t(y_1^L \cup y_2^L)$,*

$$\sum_{x \in y_1^L \cup y_2^L} f(x) = \frac{\binom{k-1}{t-1} \binom{v-1}{k-1}}{\lambda \binom{v-1}{t-1}} \sum_{x \in X} f(x).$$

Proof. (i) Let $(V, \mathcal{B})$ be a t -(v, k, λ) design, and $x \in V$. We consider the incidence structure $(V', \mathcal{B}')$, where

$$V' = V \setminus \{x\}, \quad \mathcal{B}' = \{B \in \mathcal{B} \mid x \notin B\} \cup \{B \setminus \{x\} \mid x \in B \in \mathcal{B}\}.$$

Then $(V', \mathcal{B}')$ is a regular t -wise balanced design with parameters determined by (2.4). (ii) The assertion follows by (i) and Proposition 3.1. $\square$

We close this section with some remarks on regular t -wise balanced designs. First, as far as the authors know, there are only a few general results on the existence of regular t -wise balanced designs for $t \geq 3$. Some examples are known, most of which are obtained by trivial ways as Proposition 3.4 (i). The second author and Reinhard Laue searched for regular 3-, 4- and 5-wise balanced designs with Discreta, a sophisticated program to compute designs, and found many designs with small parameters, some of which are summarized in Table 1. We believe that there will

TABLE 1. Some new regular t -wise balanced designs

Parameters	Groups
3-(25, {6, 10}, 4)	$AGL(1, 25)$
4-(27, {5, 8}, 5)	$ASL(3, 3)$
5-(33, {6, 7}, 10)	$P\Gamma L(2, 32)$
5-(33, {6, 8}, 20)	$P\Gamma L(2, 32)$
5-(33, {6, 9}, 15)	$P\Gamma L(2, 32)$
5-(33, {7, 10}, 42)	$P\Gamma L(2, 32)$
5-(55, {6, 5}, 5)	$C_2 \times P\Gamma L(2, 27)$

be further nontrivial regular t -wise balanced designs. However, in this paper, such thorough discussions are omitted and left for future work.

A natural problem is to find a good bound for the number of blocks of a t -wise balanced design. Ziqing Xiang, a student of Eiichi Bannai, recently derived the Fisher-type bound for regular t -wise balanced designs. Namely, he showed that if there is a regular $2e$ -wise balanced design $(V, \mathcal{B})$ with f distinct sizes of blocks, then

$$|\mathcal{B}| \geq \sum_{i=0}^{f-1} \binom{|V|}{e-i}.$$

This bound is sharp when $t = 2$ and $f = 2$, by a result of Woodal [36]. Moreover, when $t = 4$ and $f = 2$, a tight example can be constructed from the ordinary tight 4-design which corresponds to the Johnson scheme. Without regularity, no good bounds seem to be known ².

4. CUBATURE ARISING FROM VICTOIR'S METHOD AND ITS GENERALIZATION

In this section many cubature formulas are constructed by Victoir's method and its generalization formulated in Section 3.

²Eiichi Bannai kindly told us detailed informations on bounds for regular t -wise balanced designs through email conversation.

4.1. Index-four cubature. There are many publications on the existence of index-four cubature in small dimensional spaces that are not minimal but have few points; see, e.g., [26], [34]. In general dimensional cases, however, it seems that explicit constructions of good cubature are not enough known³. Therefore the following theorem by Shatalov [32] is very important.

Theorem 4.1. (i) ([32, Theorem 4.4.9]). *Assume that for given m, n , and q , there exists a cubature of index q with n points on S^{m-1} . Then for any $M \geq m$, there exists a cubature of index q with $((q+2)/2)^{M-m}n$ points on S^{M-1} .*

(ii) ([32, Corollary 4.4.12]). *There exists an index-four cubature on S^{m-1} with n points when*

$$(4.1) \quad m = 2^{2l} + s, \quad n = 2^{2l} \cdot 3^s \cdot (2^{2l-1} + 1), \quad l \geq 1, s \geq 0.$$

$$(4.2) \quad m = 2l + 2 + s, \quad n = 3^{s+1} \cdot ((l+1)^2 + 1), \quad l \text{ is a prime power}, s \geq 0.$$

Remark 4.2. For $s = 0$, Theorem 4.1 is a theorem of König [20]. Family (4.1) improves the upper-bound part of (2.1) if s is fixed and m is sufficiently large, or $s = 1, 2$. A similar conclusion holds for (4.2).

Cubature formulae in general-dimensional spaces that improve Shatalov's families are constructed.

Theorem 4.3. (i) *Let $l \geq 2, m$ be integers. Assume that*

$$\ell = \begin{cases} 4l - 1 & \text{if } 2^{2l-1} \leq m \leq 2^{2l}; \\ 4l + 1 & \text{if } 2^{2l} < m < 2^{2l+1}. \end{cases}$$

Then there is an integer n with $2^{\ell-1} + m < n \leq 2^\ell + m$ for which an index-four cubature with n points on S^{m-1} exists.

(ii) *Let $l \geq 2, l', m$ be integers. Assume that $m \in \{3^{l'+2} - 2, 2 \cdot 9^{l'+1} - 2\}$, and*

$$\ell = \begin{cases} 4l - 1 & \text{if } 2^{2l-1} \leq (m+2)/3 \leq 2^{2l}; \\ 4l + 1 & \text{if } 2^{2l} < (m+2)/3 < 2^{2l+1}. \end{cases}$$

Then there is an integer n with $2^{\ell-1}m < n \leq 2^\ell m$ for which an index-four cubature with n points on S^{m-1} exists.

The following lemma is employed, where the proof is easy and so omitted.

Lemma 4.4. *The following is an m -dimensional index-two cubature for $\hat{\mathcal{I}}$.*

(i) *For $m \geq 3$,*

$$\hat{\mathcal{I}}[f] = \frac{1}{2m} \sum_{x \in \mathbf{v}_1(\sqrt{4m}, 0)^L} f(x) + \frac{1}{2} \sum_{x \in \mathbf{v}_m(\sqrt{2}, 0)^L} f(x).$$

(ii) *For $m \equiv 1 \pmod{3}$,*

$$\hat{\mathcal{I}}[f] = \frac{1}{\binom{m}{(m+2)/3}} \sum_{x \in \mathbf{v}_{(m+2)/3}(\sqrt{\frac{9m}{m+2}}, 0)^L} f(x).$$

More B_m -invariant cubature can be obtained systematically by using the Sobolev theorem.

Proof of Theorem 4.3 (i). Take an $OA(2^{4l}, m, 2, 4)$ if $2^{2l-1} \leq m \leq 2^{2l}$, and an $OA(2^{4l+2}, m, 2, 4)$ if $2^{2l} < m < 2^{2l+1}$. These OA are constructed from an

³Oksana Shatalov and Yuan Xu kindly told us these informations.

$OA(2^{4l}, 2^{2l}, 2, 4)$ and an $OA(2^{4l+2}, 2^{2l+1} - 1, 2, 4)$ which are the dual of the Kerdock code and the BCH code over $\mathbb{F}_2$ (cf. [15, p. 102, p. 94]) respectively, where $0, 1 \in \mathbb{F}_2$ are replaced by $-1, 1$. Hence, by Theorem 2.12 (ii), Lemma 4.4 (i) and Proposition 2.11, we get an index-four cubature for $\mathcal{I}$ with at most $2^{\ell+1} + 2m$ points. The Kerdock OA has central symmetry (cf. [20]). The BCH OA is also centrally symmetric since it is linear. The result follows by Propositions 2.3 and 3.4.

(ii) The existence of a $2-(m, (m+2)/3, (m+2)/9)$ design with m blocks is known [17]. So, by Theorem 2.12 (i) and Lemma 4.4 (ii), we obtain an index-two cubature for $\hat{\mathcal{I}}$ with m points. According to Proposition 2.11, the resulting cubature is equivalent to an $\hat{L}$ -invariant cubature of index 4 with $2^{(m+2)/3}m$. Applying Theorem 2.12 (ii) to this formula and the OA given in the proof of Theorem 4.3, we have an index-four cubature for $\mathcal{I}$ with at most $2^{\ell+1}m$ points. Since the Kerdock and BCH OA have central symmetry, the result follows by Propositions 3.4 and 2.3. $\square$

More general-dimensional index-four cubature with $O(m^2)$ or $O(m^3)$ points can be obtained by using suitable OA, 2-designs, and regular pairwise balanced designs.

Remark 4.5. (i) Theorem 4.3 improves Theorem 4.1 for many values of m . When $2^{2l-1} \leq m \leq 2^{2l}$, the family of Theorem 4.3 (i) comes from centrally symmetric cubature by “halving” opposite row-vectors of OA. The underlying symmetric cubature were found by Victor [35, Subsection 5.3]. (ii) Theorem 4.3 does not mention the exact number of points of the constructed cubature. When $m = 2^{2l}$ in Theorem 4.3 (i), the underlying OA is the Kerdock OA and no two distinct rows coincide. So, the constructed cubature has exactly $2^{4l-1} + 2^{2l}$ points, which is equivalent to König’s family. (iii) By Proposition 2.11 the L -invariant formula of Lemma 4.4 (i) is equivalent to the degree-five cubature of Stroud [34]. Moreover the formula (ii) corresponds to Kürschák’s identity in number theory; see Section 6.

4.2. Index-six cubature. Shatalov [32, Theorem 4.7.20] compiled known index-six cubature with few points in small-dimensional spheres as Table 2 (strictly speaking, a part of the original). Nos. 1, 2, 4 are respectively in [27], [11], [9]⁴. Nos. 3, 5

TABLE 2. Index-six cubature on S^{m-1} with n points

No	1	2	3	4	5	6	7	8	9	10	11	12	13
m	3	4	5	6	7	8	9	10	11	16	17	18	23
n	11	23	41	63	113	120	480	1920	7680	2160	8640	34650	2300

are in [34], and No. 6 in [5]. To complete Table 2, Shatalov applied Theorem 4.1 (i) to one of the above formulae. For example, No. 7 has 4 times as many points as No. 6 does. According to Shatalov, Table 2 had not been updated so far, and the existence of general-dimensional index-six cubature with few points is not fully known

Two families of general-dimensional cubature that improve the upper-bound part of (2.1) are given.

Theorem 4.6. *Let Q be a prime power such that $Q \equiv 1 \pmod{6}$, $Q \neq 25$. Let $m \in \{Q + 1, Q\}$ and l be an integer with $l \geq 3$, $2^{2l-2} < m \leq 2^{2l}$. Then there is an*

⁴The existence of 23-point cubature of index 6 on S^3 is not covered in [32, Theorem 4.7.20].

integer $n \leq 2^{6l-2}(3Q(Q+1)+1)+m$ for which an index-six cubature with n points on S^{m-1} exists.

Lemma 4.7. *The following is an m -dimensional index-three cubature for $\hat{\mathcal{I}}$. (i) For $m \equiv 2 \pmod{6}$ and $8 \leq m$,*

$$\begin{aligned} \hat{\mathcal{I}}[f] = & \frac{1}{3} \sum_{x \in \mathbf{v}_m(\sqrt[3]{\frac{12}{5}}, 0)^L} f(x) + \frac{1}{3m} \sum_{x \in \mathbf{v}_1(\sqrt[3]{\frac{216m}{m+4}}, 0)^L} f(x) \\ & + \frac{1}{3(\frac{m}{(m+10)/6})} \sum_{x \in \mathbf{v}_{(m+10)/6}(\sqrt[3]{\frac{1296m(m-1)}{(m+4)(m+10)}}, 0)^L} f(x) \end{aligned}$$

(ii) For $m \equiv 1 \pmod{6}$ and $7 \leq m$,

$$\begin{aligned} \hat{\mathcal{I}}[f] = & \frac{1}{3} \sum_{x \in \mathbf{v}_m(\sqrt[3]{\frac{9}{5}}, 0)^L} f(x) + \frac{1}{3m} \sum_{x \in \mathbf{v}_1(\frac{1}{3m}, 0)^L} f(x) \\ & + \frac{1}{3(\frac{m+1}{(m+11)/6})} \sum_{x \in \mathbf{v}_{(m+11)/6}(\sqrt[3]{\frac{1296m(m+1)}{(m+5)(m+11)}}, 0)^L \cup \mathbf{v}_{(m+5)/6}(\sqrt[3]{\frac{1296m(m+1)}{(m+5)(m+11)}}, 0)^L} f(x). \end{aligned}$$

Proof of Theorem 4.6. First we consider the case where $m = Q + 1$. There exists a $3-(Q+1, (Q+11)/6, (Q+5)(Q+11)/72)$ (cf. [19]), which has $3Q(Q+1)$ blocks by (2.4). By Theorem 2.12 (i) and Lemma 4.7 (i), we obtain an index-three cubature for $\hat{\mathcal{I}}$ with $1+(Q+1)+3Q(Q+1)$ points. By Proposition 2.11, this is equivalent to an $\hat{\mathcal{I}}$ -invariant cubature with $2^{Q+1}+2(Q+1)+2^{(Q+11)/6} \cdot 3Q(Q+1)$ points. By applying Theorem 2.12 (ii) to an $OA(2^{6l-1}, Q+1, 2, 7)$ and an $OA(2^{6l-1}, (Q+11)/6, 2, 7)$ that are subarrays of the dual $OA(2^{6l-1}, 2^{2l}, 2, 7)$ of the Delsarte-Goethals code (cf. [15, p. 103]), we obtain an index-six formula for $\mathcal{I}$ with at most $2^{6l-1} \cdot (1+3Q(Q+1))+2(Q+1)$ points. Note that the $OA(2^{6l-1}, 2^{2l}, 2, 7)$ has central symmetry. In fact, the Delsarte-Goethals code can be constructed by applying the Gray-code mapping $0 \mapsto 00, 1 \mapsto 01, 2 \mapsto 11, 3 \mapsto 10$ to linear, cyclic codes over $\mathbb{Z}_4$. Replacing $0, 1 \in \mathbb{F}_2$ by ± 1 implies the central symmetry of the OA. The result thus follows by Propositions 3.4 and 2.3. Similar arguments work when $m = Q$; replace the above L -invariant formula by that of Lemma 4.7 (ii). By Proposition 3.4 (ii) the above 3-design can be reduced to a regular 3-wise balanced design with Q points and $3Q(Q+1)$ blocks. By Theorem 3.3 we obtain an index-three cubature for $\hat{\mathcal{I}}$ with $1+3Q(Q+1)+Q$ points. Then the assertion follows by the same argument as in the case $m = Q+1$. $\square$

Remark 4.8. The family of Theorem 4.6 has $O(m^5)$ points, improving the upper-bound part of (2.1). More general-dimensional index-six cubature with $O(m^5)$ points may be obtained by using known infinite families of 3-designs [19].

Two more interesting cubature are given.

Example 4.9. *The following is a 7-dimensional index-three cubature for $\hat{\mathcal{I}}$:*

$$(4.3) \quad \hat{\mathcal{I}}[f] = \frac{1}{140} \sum_{x \in \mathbf{v}_4(\sqrt[3]{28}, 0)^L \cup \mathbf{v}_3(\sqrt[3]{28}, 0)^L} f(x) + \frac{1}{14} \sum_{x \in \mathbf{v}_1(\sqrt[3]{112}, 0)^L} f(x).$$

A 3-(8, 4, 1) design exists (cf. [19]), and so does a regular 3-(7, {4, 3}, 1) design with 7 blocks of sizes 4 and 3 according to Proposition 3.4 (i). Let X be the columns of an incidence matrix of the 3-wise balanced design. By Proposition 3.4 (ii),

$$(4.4) \quad \sum_{x \in \mathbf{v}_4(\sqrt[3]{28}, 0)^L \cup \mathbf{v}_3(\sqrt[3]{28}, 0)^L} f(x) = 5 \sum_{x \in X} f(x)$$

for every $f \in \mathcal{P}_3$. Hence, by (4.3), (4.4), and Proposition 2.11, the following index-six cubature for $\mathcal{I}$ is obtained.

$$(4.5) \quad \mathcal{I}[f] = \frac{1}{448} \sum_{x \in (\sqrt[6]{28} \cdot X_1)^L} f(x) + \frac{1}{224} \sum_{x \in (\sqrt[6]{28} \cdot X_2)^L} f(x) + \frac{1}{28} \sum_{x \in \mathbf{v}_1(\sqrt[3]{112}, 0)^{B_m}} f(x)$$

where $X_1 = \{x \in X \mid \text{wt}(x) = 4\}$, $X_2 = \{x \in X \mid \text{wt}(x) = 3\}$. This is reduced to a 91-point formula of index 6 on S^6 by Propositions 2.3 and 3.4.

Example 4.10. The following is a 9-dimensional index-three cubature for $\hat{\mathcal{I}}$:

$$\hat{\mathcal{I}}[f] = \frac{1}{3} \sum_{x \in \mathbf{v}_9(1, 0)^L} f(x) + \frac{1}{630} \sum_{x \in \mathbf{v}_4(\sqrt[3]{60}, 0)^L \cup \mathbf{v}_3(\sqrt[3]{60}, 0)^L} f(x) + \frac{1}{27} \sum_{x \in \mathbf{v}_1(\sqrt[3]{180}, 0)^L} f(x).$$

The existence of a 3-(10, 4, 1) design (cf. [19]) implies that of a regular 3-(9, {4, 3}, 1) design with 12 blocks of size 3 and 18 blocks of size 4. By the same way as in Example 4.9, a 457-point formula on S^8 is obtained.

Remark 4.11. (i) The formula No. 5 of Table 2 implies that $N(7, 6) \leq 113$. Example 4.9 improves this to

$$(4.6) \quad N(7, 6) \leq 91.$$

The lower-bound part of (2.1) shows $84 \leq N(7, 6)$. The authors do not know the existence of cubature with fewer points than the 91-point formula on S^6 . It is also noted that spherical 84-point index-six cubature on S^6 do not exist by Theorem 1 of [3]. (ii) The formula No. 7 of Table 2 implies that $N(9, 6) \leq 480$. Example 4.10 improves this to

$$(4.7) \quad N(9, 6) \leq 457.$$

The fundamental roots of the group B_m are $\alpha_i = e_i - e_{i+1}$ for $i = 1, \dots, m-1$, and $\alpha_m = \sqrt{2}e_m$, where $e_1, \dots, e_m$ are the standard basis vectors in $\mathbb{R}^m$ [4]. The corner vectors are $v_i = (1/\sqrt{i}, \dots, 1/\sqrt{i}, 0, \dots, 0)$ for $i = 1, \dots, m$. We note that all B_m -invariant cubature of indices 4, 6 given in Section 4 consist of the orbits of the corner vectors. By Bajnok's theorem, in order to find higher-index spherical cubature, we must take at least one orbits of points which are not corner vectors; see, e.g., [29] for a simple construction of higher-index cubature on spheres.

A strengthening of Bajnok's theorem is proved in the next section.

5. THE MAXIMUM STRENGTH OF INVARIANT EUCLIDEAN DESIGNS

We use the notations R, J, α_i, v_i , and $\mathcal{X}(G, J)$ that are defined in Subsection 2.3. The aim of this section is to prove the following theorem.

Theorem 5.1. *Let G be a finite irreducible reflection group in $\mathbb{R}^m$ with $m \geq 2$. Then there is no choice of R, J , and a weight w for which $(\mathcal{X}(G, J), w)$ is a Euclidean t -design of $\mathbb{R}^m$ in the following cases:*

- (i) $t \geq 6$ if $G = A_{m-1}$;

- (ii) $t \geq 8$ if $G = B_m, D_m$;
- (iii) $t \geq 10$ if $G = E_6$;
- (iv) $t \geq 12$ if $G = F_4, H_3, E_7$;
- (v) $t \geq 16$ if $G = E_8$;
- (vi) $t \geq 24$ if $G = H_4$.

The following lemma plays an important role to prove the theorem.

Lemma 5.2. *Let G be a subgroup of $O(m)$, and $X = \{x_1, \dots, x_M\}$ be a subset of S^{m-1} . Let $\{f_{i,k}\}_{k=1}^{m_i}$ be a basis of $\text{Harm}_{2i}(\mathbb{R}^m)^G$, where $m_i = \dim(\text{Harm}_{2i}(\mathbb{R}^m)^G)$. Let V_i be the space $\text{Span}_{\mathbb{R}}\{(f_{i,k}(x_1), \dots, f_{i,k}(x_M)) \mid k = 1, \dots, m_i\} \subset \mathbb{R}^X$. Suppose there is $v \in \sum_{i=1}^s V_i$ such that all entries of v are positive. Then there is no choice of radii r_i and a weight w for which $(\sum_{i=1}^M r_i x_i^G, w)$ is a Euclidean $2s$ -design.*

Proof. Since $X \subset S^{m-1}$, we can express

$$\begin{aligned} v &= \sum_{i=1}^s \sum_{k=1}^{m_i} a_{i,k} (f_{i,k}(x_1), \dots, f_{i,k}(x_M)) \\ &= \sum_{i=1}^s \sum_{k=1}^{m_i} a_{i,k} (\|x_1\|_2^{2s-2i} f_{i,k}(x_1), \dots, \|x_M\|_2^{2s-2i} f_{i,k}(x_M)), \end{aligned}$$

where $a_{i,k}$ are real numbers. Let $f(x) := \sum_{i=1}^s \sum_{k=1}^{m_i} a_{i,k} \|x\|_2^{2s-2i} f_{i,k}(x)$. Then $f \in \sum_{2i+2j=2s, i \geq 1, j \geq 0} \|x\|_2^{2j} \text{Harm}_{2i}(\mathbb{R}^m)^G$, and f satisfies $f(x_i) > 0$ for each $i = 1, \dots, M$. By noting that $f(r_i x_i^g) = r_i^{2s} f(x_i^g) = r_i^{2s} f(x_i) > 0$ for $i = 1, \dots, M$, and $g \in G$, this lemma follows. $\square$

Remark 5.3. If our assumption in Lemma 5.2 holds, then any subset of $\{rx^g \mid g \in G, x \in X, r > 0\}$ does not form a Euclidean $2s$ -design. In particular, for any subgroup H of G , $(\sum_{i=1}^M r_i x_i^H, w)$ is not a Euclidean $2s$ -design for any radii r_i and weight w .


The proof of Theorem 5.1 is divided into some cases. The following notations are used. For a finite irreducible reflection group G , v_i denotes the corner vector normalized by $(v_i, \alpha_i) = 1$, $v'_i := v_i / \sqrt{(v_i, v_i)}$, and $N_i := |v_i^G|$. Let e_i be the column vector with the i -th entry 1 and the others 0. Define

$$\text{sym}(f) := \frac{1}{|(S_m)_f|} \sum_{g \in S_m} f(x^g), \quad (S_m)_f := \{g \in S_m \mid f(x^g) = f(x)\}$$

for an m -variable polynomial f , where S_m is the symmetric group of m elements. Let $p_i := x_2^2 + x_3^2 + \dots + x_{i+1}^2$ for $i \geq 2$. The polynomials h_i in the following subsections are harmonic.

5.1. Group F_4 .

Dynkin diagram



Exponents 1, 5, 7, 11.

Fundamental roots

$$\alpha_1 := {}^t e_1 - {}^t e_2, \alpha_2 := {}^t e_2 - {}^t e_3, \alpha_3 := {}^t e_4, \alpha_4 := \frac{-{}^t e_1 - {}^t e_2 - {}^t e_3 + {}^t e_4}{2}.$$

Corner Vectors

$$v_1 = {}^t e_1 + {}^t e_4, v_2 = {}^t e_1 + {}^t e_2 + 2{}^t e_4, v_3 = {}^t e_1 + {}^t e_2 + {}^t e_3 + 3 {}^t e_4, v_4 = 2{}^t e_4.$$

Size of Orbit $N_1 = 24, N_2 = 96, N_3 = 96, N_4 = 24.$

Harmonic Molien series

$$\frac{1}{(1-t^6)(1-t^8)(1-t^{12})} = 1 + t^6 + t^8 + 2t^{12} + t^{14} + \dots.$$

 G -invariant harmonic polynomials

For $i = 6, 8, 12$, $\text{Harm}_i(\mathbb{R}^4)^{F_4}$ is spanned by the following:

1. *Degree 6.*

$$f_6 := \text{sym}(x_1^6) - 5\text{sym}(x_1^4 x_2^2) + 30\text{sym}(x_1^2 x_2^2 x_3^2).$$

2. *Degree 8.*

$$f_8 := \text{sym}(x_1^8) - \frac{28}{3}\text{sym}(x_1^6 x_2^2) + \frac{98}{3}\text{sym}(x_1^4 x_2^4) - 28\text{sym}(x_1^4 x_2^2 x_3^2) + 504x_1^2 x_2^2 x_3^2 x_4^2.$$

3. *Degree 12.*

$$\begin{aligned} f_{12,1} &:= \text{sym}(x_1^{12}) - 22\text{sym}(x_1^{10} x_2^2) + 79\text{sym}(x_1^8 x_2^4) \\ &\quad + 258\text{sym}(x_1^8 x_2^2 x_3^2) - 116\text{sym}(x_1^6 x_2^6) - 236\text{sym}(x_1^6 x_2^4 x_3^2) \\ &\quad - 4392\text{sym}(x_1^6 x_2^2 x_3^2 x_4^2) + 570\text{sym}(x_1^4 x_2^4 x_3^4) + 3660\text{sym}(x_1^4 x_2^4 x_3^2 x_4^2), \end{aligned}$$

$$\begin{aligned} f_{12,2} &:= \text{sym}(x_1^{12}) - 22\text{sym}(x_1^{10} x_2^2) + \frac{133}{2}\text{sym}(x_1^8 x_2^4) \\ &\quad + \frac{591}{2}\text{sym}(x_1^8 x_2^2 x_3^2) - \frac{157}{2}\text{sym}(x_1^6 x_2^6) - \frac{1369}{4}\text{sym}(x_1^6 x_2^4 x_3^2) \\ &\quad - 4167\text{sym}(x_1^6 x_2^2 x_3^2 x_4^2) + \frac{2265}{2}\text{sym}(x_1^4 x_2^4 x_3^4) + \frac{6945}{2}\text{sym}(x_1^4 x_2^4 x_3^2 x_4^2). \end{aligned}$$

Substitute v_k for G -invariant harmonic polynomials

1. *Degree 6.*

$$u_6 := [f_6(v'_1), f_6(v'_2), f_6(v'_3), f_6(v'_4)] = [-1, -\frac{1}{9}, \frac{1}{9}, 1].$$

2. *Degree 8.*

$$u_8 := [f_8(v'_1), f_8(v'_2), f_8(v'_3), f_8(v'_4)] = [1, -\frac{13}{27}, -\frac{13}{27}, 1].$$

3. *Degree 12.*

$$u_{12,1} := [f_{12,1}(v'_1), f_{12,1}(v'_2), f_{12,1}(v'_3), f_{12,1}(v'_4)] = [0, \frac{128}{243}, -\frac{25}{243}, 1].$$


$$u_{12,2} := [f_{12,2}(v'_1), f_{12,2}(v'_2), f_{12,2}(v'_3), f_{12,2}(v'_4)] = [\frac{25}{128}, \frac{1751}{3456}, 0, 1].$$

Proposition 5.4. *There is no choice of R, J , and w for which $(\mathcal{X}(F_4, J), w)$ is a Euclidean 12-design.*

Proof. Since we have

$$-u_{12,1} + 2u_{12,2} = [\frac{25}{64}, \frac{7567}{15552}, \frac{25}{243}, 1],$$

this proposition follows by Lemma 5.2. $\square$

5.2. Group H_3 .**Dynkin diagram****Exponents** 1, 5, 9.**Fundamental roots**

$$\alpha_1 := -{}^t e_1 + {}^t e_2, \alpha_2 := -{}^t e_2 + {}^t e_3, \alpha_3 := \frac{(1+\sqrt{2}+\sqrt{5}-\sqrt{10})({}^t e_1+{}^t e_2)-(2-\sqrt{2}+2\sqrt{5}+\sqrt{10}){}^t e_3}{6}.$$

Corner Vectors

$$v_1 = \frac{-(3\sqrt{2}+\sqrt{10}+8){}^t e_1-(3\sqrt{2}+\sqrt{10}-4)({}^t e_2+{}^t e_3)}{12},$$

$$v_2 = \frac{-(3\sqrt{2}+\sqrt{10}+2)({}^t e_1+{}^t e_2)-(3\sqrt{2}+\sqrt{10}-4){}^t e_3}{6}, v_3 = -\frac{(\sqrt{2}+\sqrt{10})({}^t e_1+{}^t e_2+{}^t e_3)}{4}.$$

Size of Orbit $N_1 = 12, N_2 = 30, N_3 = 20$.**Harmonic Molien series**

$$\frac{1}{(1-t^6)(1-t^{10})} = 1 + t^6 + t^{10} + t^{12} + t^{16} + t^{18} + t^{20} + \dots.$$

 G -invariant harmonic polynomialsFor $i = 6, 10, 12$, $\text{Harm}_i(\mathbb{R}^3)^{H_3}$ is spanned by the following:1. *Degree 6.*

$$f_6 := 2\text{sym}(x_1^6) + 21\text{sym}(x_1^5 x_2) - 15\text{sym}(x_1^4 x_2^2) + 21\sqrt{10}\text{sym}(x_1^4 x_2 x_3) \\ - (70 - 7\sqrt{10})\text{sym}(x_1^3 x_2^3) - 21\sqrt{10}\text{sym}(x_1^3 x_2^2 x_3) + 180x_1^2 x_2^2 x_3^2.$$

2. *Degree 10.*

$$f_{10} := \sum_{g \in H_3} h_{10}(x^g),$$

where

$$h_{10}(x) := 256x_1^{10} - 5760x_1^8 p_2 + 20160x_1^6 p_2^2 - 16800x_1^4 p_2^3 + 3150x_1^2 p_2^4 - 63p_2^5.$$

3. *Degree 12.*

$$f_{12} := \sum_{g \in H_3} h_{12}(x^g),$$

where

$$h_{12}(x) := 1024x_1^{12} - 33792x_1^{10} p_2 + 190080x_1^8 p_2^2 \\ - 295680x_1^6 p_2^3 + 138600x_1^4 p_2^4 - 16632x_1^2 p_2^5 + 231p_2^6.$$

Substitute v_k for G -invariant harmonic polynomials1. *Degree 6.*

$$u_6 := [f_6(v'_1), f_6(v'_2), f_6(v'_3)] = [\frac{14\sqrt{10}-4}{5}, \frac{-7\sqrt{10}+2}{8}, \frac{-14\sqrt{10}+4}{9}].$$

2. *Degree 10.*

$$u_{10} := [f_{10}(v'_1), f_{10}(v'_2), f_{10}(v'_3)] \\ = [-\frac{43124224\sqrt{10}+49637120}{98415}, \frac{8422700\sqrt{10}+9694750}{19683}, -\frac{1078105600\sqrt{10}+1240928000}{1594323}].$$

3. *Degree 12.*


$$\begin{aligned} u_{12} &:= [f_{12}(v'_1), f_{12}(v'_2), f_{12}(v'_3)] \\ &= \left[\frac{191679488\sqrt{10}-6897476096}{492075}, \frac{10856846\sqrt{10}-390677357}{39366}, \frac{-191679488\sqrt{10}+6897476096}{14348907} \right]. \end{aligned}$$

Proposition 5.5. *There is no choice of R, J , and w for which $(\mathcal{X}(H_3, J), w)$ is a Euclidean 12-design.*

Proof. There is $u \in \text{Span}_{\mathbb{R}}\{u_6, u_{10}, u_{12}\}$ all whose entries are positive, since the vectors u_6, u_{10}, u_{12} are linearly independent. The result follows by Lemma 5.2. $\square$

5.3. **Group H_4 .**

Dynkin diagram



Exponents 1, 11, 19, 29.

Fundamental roots

$$\alpha_1 := -{}^t e_1 + {}^t e_2, \alpha_2 := -{}^t e_2 + {}^t e_3, \alpha_3 := -{}^t e_3 + {}^t e_4, \alpha_4 := \frac{{}^t e_1 + {}^t e_2 + {}^t e_3 + \sqrt{5} {}^t e_4}{2}.$$

Corner Vectors

$$\begin{aligned} v_1 &= \frac{(\sqrt{5}-1){}^t e_1 + (\sqrt{5}+3)({}^t e_2 + {}^t e_3 - {}^t e_4)}{4}, v_2 = \frac{(\sqrt{5}+1)({}^t e_1 + {}^t e_2) + (\sqrt{5}+3)({}^t e_3 - {}^t e_4)}{2}, \\ v_3 &= \frac{(3\sqrt{5}+5)({}^t e_1 + {}^t e_2 + {}^t e_3) - 3(\sqrt{5}+3){}^t e_4}{4}, v_4 = \frac{(\sqrt{5}+3)({}^t e_1 + {}^t e_2 + {}^t e_3 - {}^t e_4)}{2}. \end{aligned}$$

Size of Orbit $N_1 = 120, N_2 = 720, N_3 = 1200, N_4 = 600$.

Harmonic Molien series

$$\frac{1}{(1-t^{12})(1-t^{20})(1-t^{30})} = 1 + t^{12} + t^{20} + t^{24} + t^{30} + \dots$$

G -invariant harmonic polynomials

For $i = 12, 20, 24$, $\text{Harm}_i(\mathbb{R}^4)^{H_4}$ is spanned by the following:

1. *Degree 12.*

$$f_{12} := \sum_{g \in H_4} h_{12}(x^g),$$

where

$$h_{12}(x) := 13x_1^{12} - 286x_1^{10}p_3 + 1287x_1^8p_3^2 - 1716x_1^6p_3^3 + 715x_1^4p_3^4 - 78x_1^2p_3^5 + p_3^6.$$

2. *Degree 20.*

$$f_{20} := \sum_{g \in H_4} h_{20}(x^g),$$

where

$$\begin{aligned} h_{20}(x) &:= 21x_1^{20} - 1330x_1^{18}p_3 + 20349x_1^{16}p_3^2 - 116280x_1^{14}p_3^3 + 293930x_1^{12}p_3^4 \\ &\quad - 352716x_1^{10}p_3^5 + 203490x_1^8p_3^6 - 54264x_1^6p_3^7 + 5985x_1^4p_3^8 - 210x_1^2p_3^9 + p_3^{10}. \end{aligned}$$

3. *Degree 24.*

$$f_{24} := \sum_{g \in H_4} h_{24}(x^g),$$

where

$$\begin{aligned} h_{24}(x) := & x_1^{24} - 92x_1^{22}p_3 + \frac{10626}{5}x_1^{20}p_3^2 - 19228x_1^{18}p_3^3 + 81719x_1^{16}p_3^4 \\ & - 178296x_1^{14}p_3^5 + 208012x_1^{12}p_3^6 - \frac{653752}{5}x_1^{10}p_3^7 + 43263x_1^8p_3^8 \\ & - 7084x_1^6p_3^9 + 506x_1^4p_3^{10} - 12x_1^2p_3^{11} + \frac{1}{25}p_3^{12}. \end{aligned}$$

Substitute v_k for G -invariant harmonic polynomials

1. *Degree 12.*

$$u_{12} := [f_{12}(v'_1), f_{12}(v'_2), f_{12}(v'_3), f_{12}(v'_4)] = [-4500, 540, \frac{32500}{27}, \frac{5625}{4}].$$

2. *Degree 20.*

$$u_{20} := [f_{20}(v'_1), f_{20}(v'_2), f_{20}(v'_3), f_{20}(v'_4)] = [6975, -\frac{58869}{25}, \frac{4035425}{2187}, \frac{216225}{64}].$$

3. *Degree 24.*

$$u_{24} := [f_{24}(v'_1), f_{24}(v'_2), f_{24}(v'_3), f_{24}(v'_4)] = [-\frac{2367}{16}, -\frac{4689027}{50000}, \frac{416329}{104976}, \frac{622521}{16384}].$$

Proposition 5.6. *There is no choice of R, J , and w for which $(\mathcal{X}(H_4, J), w)$ is a Euclidean 24-design.*


Proof. Since we have

$$u_{20} - 30u_{24} = [\frac{91305}{8}, \frac{2293281}{5000}, \frac{30201755}{17496}, \frac{18338985}{8192}],$$

this proposition follows by Lemma 5.2. $\square$

5.4. Group E_6 .

Dynkin diagram



Exponents 1, 4, 5, 7, 8, 11.

Fundamental roots

$$\alpha_1 := {}^te_1 - {}^te_2, \alpha_2 := {}^te_2 - {}^te_3, \alpha_3 := {}^te_3 - {}^te_4, \alpha_4 := {}^te_4 - {}^te_5, \alpha_5 := {}^te_5 - {}^te_6,$$

$$\alpha_6 := \frac{(-3+\sqrt{3})({}^te_1+{}^te_2+{}^te_3)+(3+\sqrt{3})({}^te_4+{}^te_5+{}^te_6)}{6}.$$

Corner Vectors

$$\begin{aligned} v_1 &= \frac{(\sqrt{3}+5){}^te_1+(\sqrt{3}-1)({}^te_2+{}^te_3+{}^te_4+{}^te_5+{}^te_6)}{6}, v_2 = \frac{(\sqrt{3}+2)({}^te_1+{}^te_2)+(\sqrt{3}-1)({}^te_3+{}^te_4+{}^te_5+{}^te_6)}{3}, \\ v_3 &= \frac{(\sqrt{3}+1)({}^te_1+{}^te_2+{}^te_3)+(\sqrt{3}-1)({}^te_4+{}^te_5+{}^te_6)}{2}, v_4 = \frac{(\sqrt{3}+1)({}^te_1+{}^te_2+{}^te_3+{}^te_4)+(\sqrt{3}-2)({}^te_5+{}^te_6)}{3}, \\ v_5 &= \frac{(\sqrt{3}+1)({}^te_1+{}^te_2+{}^te_3+{}^te_4+{}^te_5)+(\sqrt{3}-5){}^te_6}{6}, v_6 = \frac{\sqrt{3}({}^te_1+{}^te_2+{}^te_3+{}^te_4+{}^te_5+{}^te_6)}{3}. \end{aligned}$$

Size of Orbit $N_1 = 27, N_2 = 216, N_3 = 720, N_4 = 216, N_5 = 27, N_6 = 72.$

Harmonic Molien series

$$\frac{1}{(1-t^5)(1-t^6)(1-t^8)(1-t^9)(1-t^{12})} = 1 + t^5 + t^6 + t^8 + t^9 + t^{10} + \dots$$

G -invariant harmonic polynomials

For $i = 5, 6, 8, 9, 10$, $\text{Harm}_i(\mathbb{R}^6)^{E_6}$ is spanned by the following:

1. *Degree 5.*

$$f_5 := \text{sym}(x_1^5) + \text{sym}(x_1^4 x_2) - 2\text{sym}(x_1^3 x_2^2) + \text{sym}(x_1^3 x_2 x_3) \\ - 3\text{sym}(x_1^2 x_2 x_3 x_4) + 24\text{sym}(x_1 x_2 x_3 x_4 x_5).$$

2. *Degree 6.*

$$f_6 := \text{sym}(x_1^6) + \frac{3}{2}\text{sym}(x_1^5 x_2) + -3\text{sym}(x_1^4 x_2^2) + \frac{15}{14}\text{sym}(x_1^4 x_2 x_3) \\ + \frac{5}{7}\text{sym}(x_1^3 x_2^3) - \frac{30}{7}\text{sym}(x_1^3 x_2^2 x_3) + \frac{30}{7}\text{sym}(x_1^3 x_2 x_3 x_4) + 9\text{sym}(x_1^2 x_2^2 x_3^2) \\ + \frac{45}{7}\text{sym}(x_1^2 x_2^2 x_3 x_4) - \frac{180}{7}\text{sym}(x_1^2 x_2 x_3 x_4 x_5) + \frac{180}{7}x_1 x_2 x_3 x_4 x_5 x_6.$$

3. *Degree 8.*

$$f_8 := \sum_{g \in E_6} h_8(x^g),$$

where

$$h_8(x) := x_1^8 - \frac{28}{5}x_1^6 p_5 + 6x_1^4 p_5^2 - \frac{4}{3}x_1^2 p_5^3 + \frac{1}{33}p_5^4.$$

4. *Degree 9.*

$$f_9 := \sum_{g \in E_6} h_9(x^g),$$

where

$$h_9(x) := \text{sym}(x_1^9) - \frac{36}{5}\text{sym}(x_1^7 x_2^2) + \frac{126}{5}\text{sym}(x_1^5 x_2^4) - 63\text{sym}(x_1^4 x_2^3 x_3^2) \\ + 63\text{sym}(x_1^4 x_2^2 x_3^2 x_4) + 252\text{sym}(x_1^3 x_2^2 x_3^2 x_4^2) - 945\text{sym}(x_1^2 x_2^2 x_3^2 x_4^2 x_5).$$

5. *Degree 10.*

$$f_{10} := \sum_{g \in E_6} h_{10}(x^g),$$

where

$$h_{10}(x) := x_1^{10} - 9x_1^8 p_5 + 18x_1^6 p_5^2 - 10x_1^4 p_5^3 + \frac{15}{11}x_1^2 p_5^4 - \frac{3}{143}p_5^5.$$

Substitute v_k for G -invariant harmonic polynomials

1. *Degree 5.*

$$u_5 := [f_5(v'_1), f_5(v'_2), f_5(v'_3), f_5(v'_4), f_5(v'_5), f_5(v'_6)] \\ = [\frac{3\sqrt{3}}{4}, \frac{6\sqrt{30}}{125}, 0, -\frac{6\sqrt{30}}{125}, -\frac{3\sqrt{3}}{4}, 0].$$

2. *Degree 6.*

$$u_6 := [f_6(v'_1), f_6(v'_2), f_6(v'_3), f_6(v'_4), f_6(v'_5), f_6(v'_6)] \\ = [\frac{81}{56}, -\frac{81}{700}, -\frac{9}{28}, -\frac{81}{700}, \frac{81}{56}, -\frac{27}{28}].$$

3. *Degree 8.*

$$u_8 := [f_8(v'_1), f_8(v'_2), f_8(v'_3), f_8(v'_4), f_8(v'_5), f_8(v'_6)] \\ = [800, -\frac{6784}{25}, -\frac{640}{9}, -\frac{6784}{25}, 800, \frac{3200}{3}].$$

4. Degree 9.

$$\begin{aligned} u_9 &:= [f_9(v'_1), f_9(v'_2), f_9(v'_3), f_9(v'_4), f_9(v'_5), f_9(v'_6)] \\ &= [2065\sqrt{3}, -\frac{185024\sqrt{30}}{625}, 0, \frac{185024\sqrt{30}}{625}, -2065\sqrt{3}, 0]. \end{aligned}$$

5. Degree 10.

$$\begin{aligned} u_{10} &:= [f_{10}(v'_1), f_{10}(v'_2), f_{10}(v'_3), f_{10}(v'_4), f_{10}(v'_5), f_{10}(v'_6)] \\ &= [\frac{11520}{13}, \frac{423936}{1625}, \frac{51200}{351}, \frac{423936}{1625}, \frac{11520}{13}, -\frac{10240}{39}]. \end{aligned}$$

Proposition 5.7. *There is no choice of R, J , and w for which $(\mathcal{X}(E_6, J), w)$ is a Euclidean 10-design.*


Proof. Since we have

$$u_{10} + u_8 = [\frac{11745}{2816}, \frac{13527}{220000}, \frac{387}{1760}, \frac{13527}{220000}, \frac{11745}{2816}, \frac{621}{352}],$$

this proposition follows by Lemma 5.2. $\square$

5.5. Group E_7 .

Dynkin diagram



Exponents 1, 5, 7, 9, 11, 13, 17.

Fundamental roots

$$\begin{aligned} \alpha_1 &:= {}^t e_1 - {}^t e_2, \alpha_2 := {}^t e_2 - {}^t e_3, \alpha_3 := {}^t e_3 - {}^t e_4, \alpha_4 := {}^t e_4 - {}^t e_5, \alpha_5 := {}^t e_5 - {}^t e_6, \\ \alpha_6 &:= {}^t e_6 - {}^t e_7, \alpha_7 := \frac{(-4+\sqrt{2})({}^t e_1+{}^t e_2+{}^t e_3)+(3+\sqrt{2})({}^t e_4+{}^t e_5+{}^t e_6+{}^t e_7)}{7}. \end{aligned}$$

Corner Vectors

$$\begin{aligned} v_1 &= \frac{(6+2\sqrt{2}){}^t e_1+(-1+2\sqrt{2})({}^t e_2+{}^t e_3+{}^t e_4+{}^t e_5+{}^t e_6+{}^t e_7)}{7}, \\ v_2 &= \frac{(5+4\sqrt{2})({}^t e_1+{}^t e_2)+(-2+4\sqrt{2})({}^t e_3+{}^t e_4+{}^t e_5+{}^t e_6+{}^t e_7)}{7}, \\ v_3 &= \frac{(4+6\sqrt{2})({}^t e_1+{}^t e_2+{}^t e_3)+(-3+6\sqrt{2})({}^t e_4+{}^t e_5+{}^t e_6+{}^t e_7)}{7}, \\ v_4 &= \frac{(6+9\sqrt{2})({}^t e_1+{}^t e_2+{}^t e_3+{}^t e_4)+(-8+9\sqrt{2})({}^t e_5+{}^t e_6+{}^t e_7)}{14}, \\ v_5 &= \frac{(2+3\sqrt{2})({}^t e_1+{}^t e_2+{}^t e_3+{}^t e_4+{}^t e_5)+(-5+3\sqrt{2})({}^t e_6+{}^t e_7)}{7}, \\ v_6 &= \frac{(-2-3\sqrt{2})({}^t e_1+{}^t e_2+{}^t e_3+{}^t e_4+{}^t e_5+{}^t e_6)+(12-3\sqrt{2}){}^t e_7}{14}, \\ v_7 &= \frac{{}^t e_1+{}^t e_2+{}^t e_3+{}^t e_4+{}^t e_5+{}^t e_6+{}^t e_7}{\sqrt{2}}. \end{aligned}$$

Size of Orbit

$$N_1 = 126, N_2 = 2016, N_3 = 10080, N_4 = 4032, N_5 = 756, N_6 = 56, N_7 = 576.$$

Harmonic Molien series

$$\frac{1}{(1-t^6)(1-t^8)(1-t^{10})(1-t^{12})(1-t^{14})(1-t^{18})} = 1 + t^6 + t^8 + t^{10} + 2t^{12} + \dots$$

G -invariant harmonic polynomials

For $i = 6, 8, 10, 12$, $\text{Harm}_i(\mathbb{R}^7)^{E_7}$ is spanned by the following:

1. *Degree 6.*

$$f_6 := \sum_{g \in E_7} h_6(x^g),$$

where

$$h_6(x) := 32x_1^6 - 80x_1^4p_6 + 30x_1^2p_6^2 - p_6^3.$$

2. *Degree 8.*

$$f_8 := \sum_{g \in E_7} h_8(x^g),$$

where

$$h_8(x) := 384x_1^8 - 1792x_1^6p_6 + 1680x_1^4p_6^2 - 336x_1^2p_6^3 + 7p_6^4.$$

3. *Degree 10.*

$$f_{10} := \sum_{g \in E_7} h_{10}(x^g),$$

where

$$h_{10}(x) := 256x_1^{10} - 1920x_1^8p_6 + 3360x_1^6p_6^2 - 1680x_1^4p_6^3 + 210x_1^2p_6^4 - 3p_6^5.$$

4. *Degree 12.*

$$f_{12,1} := \sum_{g \in E_7} h_{12,1}(x^g), \quad f_{12,2} := \sum_{g \in E_7} h_{12,2}(x^g),$$

where

$$\begin{aligned} h_{12,1}(x) &:= 4096x_1^{12} - 45056x_1^{10}p_6 + 126720x_1^8p_6^2 \\ &\quad - 118272x_1^6p_6^3 + 36960x_1^4p_6^4 - 3168x_1^2p_6^5 + 33p_6^6, \end{aligned}$$

$$h_{12,2}(x) := x_1x_2(2048x_1^{10} - 14080x_1^8p_6 + 25344x_1^6p_6^2 - 14784x_1^4p_6^3 + 2640x_1^2p_6^4 - 99p_6^5).$$

Substitute v_k for G -invariant harmonic polynomials

1. *Degree 6.*

$$\begin{aligned} u_6 &:= [f_6(v'_1), f_6(v'_2), f_6(v'_3), f_6(v'_4), f_6(v'_5), f_6(v'_6), f_6(v'_7)] \\ &= \left[\frac{-7700659200 + 9488793600\sqrt{2}}{16807}, \frac{-427814400 + 527155200\sqrt{2}}{2401}, \frac{-1818211200 + 2240409600\sqrt{2}}{16807}, \right. \\ &\quad \left. \frac{-547602432 + 674758656\sqrt{2}}{16807}, \frac{2887747200 - 3558297600\sqrt{2}}{16807}, \frac{20535091200 - 25303449600\sqrt{2}}{16807}, \right. \\ &\quad \left. \frac{-123210547200 + 151820697600\sqrt{2}}{823543} \right]. \end{aligned}$$

2. *Degree 8.*

$$\begin{aligned} u_8 &:= [f_8(v'_1), f_8(v'_2), f_8(v'_3), f_8(v'_4), f_8(v'_5), f_8(v'_6), f_8(v'_7)] \\ &= \left[\frac{6579988992000 - 5480856576000\sqrt{2}}{823543}, \frac{731109888000 - 608984064000\sqrt{2}}{823543}, \right. \\ &\quad \left. \frac{-3527605209600 + 2938348108800\sqrt{2}}{823543}, \frac{-3134999199744 + 2611323666432\sqrt{2}}{823543}, \right. \\ &\quad \left. \frac{-1809496972800 + 1507235558400\sqrt{2}}{823543}, \frac{3509327462400 - 2923123507200\sqrt{2}}{823543}, \right. \\ &\quad \left. \frac{-115807806259200 + 96463075737600\sqrt{2}}{40353607} \right]. \end{aligned}$$

3. Degree 10.

$$\begin{aligned}
u_{10} &:= [f_{10}(v'_1), f_{10}(v'_2), f_{10}(v'_3), f_{10}(v'_4), f_{10}(v'_5), f_{10}(v'_6), f_{10}(v'_7)] \\
&= \left[\frac{-6428624451840+415928908800\sqrt{2}}{5764801}, \frac{357145802880-23107161600\sqrt{2}}{823543}, \right. \\
&\quad \frac{2388412556760-154529143200\sqrt{2}}{5764801}, \frac{-73143460429824+4732346695680\sqrt{2}}{720600125}, \\
&\quad \frac{-7433097022440+480917800800\sqrt{2}}{5764801}, \frac{30476441845760-1971811123200\sqrt{2}}{5764801}, \\
&\quad \left. \frac{3291455719342080-212955601305600\sqrt{2}}{1977326743} \right].
\end{aligned}$$

4. Degree 12.

$$\begin{aligned}
u_{12,1} &:= [f_{12,1}(v'_1), f_{12,1}(v'_2), f_{12,1}(v'_3), f_{12,1}(v'_4), f_{12,1}(v'_5), f_{12,1}(v'_6), f_{12,1}(v'_7)] \\
&= \left[\frac{27363005574796800+17942314142016000\sqrt{2}}{1977326743}, \frac{-760132890073600-689327933184000\sqrt{2}}{282475249}, \right. \\
&\quad \frac{-513174301527400-792264524693400\sqrt{2}}{1977326743}, \frac{-14026148038967296-72141536776421376\sqrt{2}}{49433168575}, \\
&\quad \frac{3931481294451000-4960153279164600\sqrt{2}}{1977326743}, \frac{-12979679661260800-37832882828083200\sqrt{2}}{1977326743}, \\
&\quad \left. \frac{249757080640811827200+284898146732782387200\sqrt{2}}{33232930569601} \right].
\end{aligned}$$

$$\begin{aligned}
u_{12,2} &:= [f_{12,2}(v'_1), f_{12,2}(v'_2), f_{12,2}(v'_3), f_{12,2}(v'_4), f_{12,2}(v'_5), f_{12,2}(v'_6), f_{12,2}(v'_7)] \\
&= \left[\frac{-2419675164360000-1489162193640000\sqrt{2}}{1977326743}, \frac{113867977056000+18727597152000\sqrt{2}}{282475249}, \right. \\
&\quad \frac{156757191916575-26126480038725\sqrt{2}}{1977326743}, \frac{16622260339703808-6701937797136384\sqrt{2}}{49433168575}, \\
&\quad \frac{1494452243214675-1107985853945025\sqrt{2}}{1977326743}, \frac{8313279170969600-2771226582220800\sqrt{2}}{1977326743}, \\
&\quad \left. \frac{-51686387833407897600+773622407142604800\sqrt{2}}{33232930569601} \right].
\end{aligned}$$

Proposition 5.8. *There is no choice of R, J , and w for which $(\mathcal{X}(E_7, J), w)$ is a Euclidean 12-design.*


Proof. Since we have

$$\begin{aligned}
&-2u_{12,1} - 25u_{12,2} + u_{10} = \\
&[2.86443 \times 10^6, 256489., 513956., 989894., 2.86352 \times 10^6, 1.64917 \times 10^7, 293023.],
\end{aligned}$$

this proposition follows by Lemma 5.2. $\square$

5.6. Group E_8 .

Dynkin diagram



Exponents 1, 7, 11, 13, 17, 19, 23, 29.

Fundamental roots

$$\begin{aligned}
\alpha_1 &:= {}^t e_1 - {}^t e_2, \alpha_2 := {}^t e_2 - {}^t e_3, \alpha_3 := {}^t e_3 - {}^t e_4, \alpha_4 := {}^t e_4 - {}^t e_5, \alpha_5 := {}^t e_5 - {}^t e_6, \\
\alpha_6 &:= {}^t e_6 - {}^t e_7, \alpha_7 := {}^t e_7 - {}^t e_8, \alpha_8 := \frac{-{}^t e_1 - {}^t e_2 - {}^t e_3 + {}^t e_4 + {}^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8}{2}.
\end{aligned}$$

Corner Vectors

$$\begin{aligned}
v_1 &= \frac{3^t e_1 + {}^t e_2 + {}^t e_3 + {}^t e_4 + {}^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8}{2}, \\
v_2 &= 2^t e_1 + 2^t e_2 + {}^t e_3 + {}^t e_4 + {}^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8, \\
v_3 &= \frac{5^t e_1 + 5^t e_2 + 5^t e_3 + 3^t e_4 + 3^t e_5 + 3^t e_6 + 3^t e_7 + 3^t e_8}{2}, \\
v_4 &= 2^t e_1 + 2^t e_2 + 2^t e_3 + 2^t e_4 + {}^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8, \\
v_5 &= \frac{3^t e_1 + 3^t e_2 + 3^t e_3 + 3^t e_4 + 3^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8}{2}, \\
v_6 &= -{}^t e_1 - {}^t e_2 - {}^t e_3 - {}^t e_4 - {}^t e_5 - {}^t e_6, \\
v_7 &= \frac{-{}^t e_1 - {}^t e_2 - {}^t e_3 - {}^t e_4 - {}^t e_5 - {}^t e_6 - {}^t e_7 + {}^t e_8}{2}, \\
v_8 &= {}^t e_1 + {}^t e_2 + {}^t e_3 + {}^t e_4 + {}^t e_5 + {}^t e_6 + {}^t e_7 + {}^t e_8.
\end{aligned}$$

Size of Orbit

$$N_1 = 2160, N_2 = 69120, N_3 = 483840, N_4 = 241920,$$

$$N_5 = 60480, N_6 = 6720, N_7 = 240, N_8 = 17280.$$

Harmonic Molien series

$$\frac{1}{(1-t^8)(1-t^{12})(1-t^{14})(1-t^{18})(1-t^{20})(1-t^{24})(1-t^{30})} = \\
1 + t^8 + t^{12} + t^{14} + t^{16} + t^{18} + 2t^{20} + \dots$$

 G -invariant harmonic polynomials

For $i = 8, 12, 14, 16$, $\text{Harm}_8(\mathbb{R}^8)^{E_8}$ is spanned by the following:

1. *Degree 8.*

$$f_8 := \sum_{g \in E_8} h_8(x^g),$$

where

$$h_8(x) := 429x_1^8 - 1716x_1^6 p_7 + 1430x_1^4 p_7^2 - 260x_1^2 p_7^3 + 5p_7^4.$$

2. *Degree 12.*

$$f_{12} := \sum_{g \in E_8} h_{12}(x^g),$$

where

$$h_{12}(x) := 1547x_1^{12} - 14586x_1^{10} p_7 + 36465x_1^8 p_7^2 - 30940x_1^6 p_7^3 + 8925x_1^4 p_7^4 - 714x_1^2 p_7^5 + 7p_7^6.$$

3. *Degree 14.*

$$f_{14} := \sum_{g \in E_8} h_{14}(x^g),$$

where

$$\begin{aligned}
h_{14}(x) &:= 969x_1^{14} - 12597x_1^{12} p_7 + 46189x_1^{10} p_7^2 - 62985x_1^8 p_7^3 \\
&\quad + 33915x_1^6 p_7^4 - 6783x_1^4 p_7^5 + 399x_1^2 p_7^6 - 3p_7^7.
\end{aligned}$$

4. *Degree 16.*

$$f_{16} := \sum_{g \in E_8} h_{16}(x^g),$$

where

$$h_{16}(x) := 6783x_1^{16} - 116280x_1^{14}p_7 + 587860x_1^{12}p_7^2 - 1175720x_1^{10}p_7^3 \\ + 1017450x_1^8p_7^4 - 379848x_1^6p_7^5 + 55860x_1^4p_7^6 - 2520x_1^2p_7^7 + 15p_7^8.$$

Substitute v_k for G -invariant harmonic polynomials

1. *Degree 8.*

$$u_8 := [f_8(v'_1), f_8(v'_2), f_8(v'_3), f_8(v'_4), f_8(v'_5), f_8(v'_6), f_8(v'_7), f_8(v'_8)] \\ = [174182400, \frac{4926873600}{49}, 82059264, 62705664, \\ 19353600, -116121600, -1045094400, 97977600].$$

2. *Degree 12.*

$$u_{12} := [f_{12}(v'_1), f_{12}(v'_2), f_{12}(v'_3), f_{12}(v'_4), f_{12}(v'_5), f_{12}(v'_6), f_{12}(v'_7), f_{12}(v'_8)] \\ = [1680315840, \frac{15655887360}{49}, \frac{14950365696}{125}, -\frac{2608490304}{125}, \\ -275607360, -734952960, 4480842240, 148777965].$$

3. *Degree 14.*

$$u_{14} := [f_{14}(v'_1), f_{14}(v'_2), f_{14}(v'_3), f_{14}(v'_4), f_{14}(v'_5), f_{14}(v'_6), f_{14}(v'_7), f_{14}(v'_8)] \\ = [1207483200, -\frac{567924825600}{16807}, -\frac{2009165312}{15}, -\frac{671799744}{5}, \\ -\frac{253422400}{3}, 184307200, -2634508800, -293294925].$$

4. *Degree 16.*

$$u_{16} := [f_{16}(v'_1), f_{16}(v'_2), f_{16}(v'_3), f_{16}(v'_4), f_{16}(v'_5), f_{16}(v'_6), f_{16}(v'_7), f_{16}(v'_8)] \\ = [1490121360, -\frac{393199971840}{2401}, \frac{3287394820358656}{1265625}, \frac{36512571016971}{62500}, \\ \frac{1232569520}{3}, 2075906560, 7529034240, -\frac{9749511135}{16}].$$

Proposition 5.9. *There is no choice of R, J , and w for which $(\mathcal{X}(E_8, J), w)$ is a Euclidean 16-design.*

Proof. Since we have

$$u_{16} - 3u_{14} + 2u_{12} = [1228303440, \frac{9691313402880}{16807}, \frac{4098709695302656}{1265625}, \frac{59096571112971}{62500}, \\ \frac{339192560}{3}, 53079040, 24394245120, \frac{9089540145}{16}],$$

this proposition follows by Lemma 5.2. $\square$

Now, we are ready to complete the proof of Theorem 5.1.

Proof of Theorem 5.1: The case (1) is in Theorem 2.10, and the cases (2), (3) in [24]. Thus the theorem follows by Propositions 5.4-5.9. $\square$

The following result, together with Theorem 5.1, determine the maximum degree of spherical cubature formulae $(\mathcal{X}(G, J), w)$ for all irreducible reflection groups G .

Theorem 5.10. *(i) An F_4 -invariant cubature of degree 11 that consists of the orbits of the corner vectors is classified by:*

$$w_1 = \frac{13-960w_4}{960}, \quad w_2 = \frac{3(-1+192w_4)}{256}, \quad w_3 = \frac{3(1-120w_4)}{160}, \quad \frac{1}{192} \leq w_4 \leq \frac{1}{120}.$$

(ii) An H_3 -invariant cubature of degree 11 that consists of the orbits of the corner vectors is classified by:

$$w_1 = \frac{125}{5544}, \quad w_2 = \frac{64}{3465}, \quad w_3 = \frac{27}{3080}.$$

(iii) An H_4 -invariant cubature of degree 23 that consists of the orbits of the corner vectors is classified by:

$$w_1 = \frac{368-9625w_4}{315392}, \quad w_2 = \frac{125(16+5625w_4)}{2359296}, \quad w_3 = -\frac{6561(16-51975w_4)}{504627200}, \quad 0 \leq w_4 \leq \frac{16}{51975}.$$

(iv) An E_6 -invariant cubature of degree 9 that consists of the orbits of the corner vectors is classified by:

$$w_1 = \frac{2(1-96w_6)}{729}, \quad w_2 = \frac{125(1+1200w_6)}{186624}, \quad w_3 = \frac{1}{1280} - \frac{9w_6}{16}, \\ w_4 = \frac{125(1+1200w_6)}{186624}, \quad w_5 = \frac{2(1-96w_6)}{729}, \quad 0 \leq w_6 \leq \frac{1}{720}.$$

(v) An E_7 -invariant cubature of degree 11 that consists of the orbits of the corner vectors is classified by the following two types of weights:

$$(1) \quad w_1 = -\frac{4(-296924467+966078461040w_2+107900687895w_3+95875084800w_7)}{610410794301}, \\ w_4 = -\frac{625(-945994+3215011030w_2+24066363475w_3+1769169600w_7)}{4340698981696}, \\ w_5 = \frac{8(34900936+247702641648w_2+1231161574335w_3+182083866624w_7)}{1831232382903}, \\ w_6 = -\frac{27(-32430307+60983896974w_2+30607311735w_3+25518620160w_7)}{542587372712}, \\ 0 \leq w_7 \leq -\frac{2401(-394+1339030w_2+10023475w_3)}{1769169600}, \quad 0 \leq w_3 < -\frac{2(-197+669515w_2)}{10023475}, \\ 0 \leq w_2 \leq \frac{197}{669515}. \\ (2) \quad w_1 = -\frac{4(-211+686070w_2)}{440055}, \quad w_3 = -\frac{2(-197+669515w_2)}{10023475}, \quad w_5 = \frac{16(1231+1230075w_2)}{54126765}, \\ w_6 = -\frac{351(-71+129360w_2)}{16037560}, \quad 0 \leq w_2 \leq \frac{197}{669515}, \quad w_4 = 0, \quad w_7 = 0.$$

(vi) An E_8 -invariant cubature of degree 15 that consists of the orbits of the corner vectors is classified by the nonnegative solutions w_i of the system of equations

$$(5.1) \quad u_8^t v = 0, \quad u_{12}^t v = 0, \quad u_{14}^t v = 0, \quad \sum_{i=1}^8 N_i w_i = 1,$$

where $v = (N_1 w_1, \dots, N_8 w_8)$, and u_i , N_i are defined in Subsection 5.6. The precise solutions of (5.1) are referred to Appendix.

Remark 5.11. The H_3 -invariant cubature of Theorem 5.10 (i) was constructed by Goethals and Seidel [9, p. 214] who found, moreover, a spherical cubature of degree 15 by taking the orbits of v'_1, v'_2, v'_3 , plus one more orbit; for example, see [12] for further informations on the existence of three-dimensional spherical cubature. It is also interesting to note that the formula given in Theorem 5.10 (vi) is equivalent to a 26400-point cubature of degree 15 which comes from shells of the Korkin-Zorotalev lattice [13]. In [9, p. 214], Goethals and Seidel found a spherical cubature of degree 19 that consists of the H_4 -orbits of the zeros of an invariant harmonic homogeneous polynomial of degree 12. Salihov [28] found another H_4 -invariant cubature of degree 19 by taking the union of the 120-cell and the 600-cell. Motivated by this, the authors searched three and four H_4 -orbits of the corner vectors, and found the higher-degree cubature of Theorem 5.10 (ii).

6. HILBERT IDENTITIES AND CUBATURE FORMULAE

As explained in Section 2, there is a cubature of index q on S^{m-1} with n points if and only if there are n vectors $r_1, \dots, r_n \in \mathbb{R}^m$ such that

$$(6.1) \quad \sum_{i=1}^n \langle x, r_i \rangle^q = \langle x, x \rangle^{\frac{q}{2}}$$

for every $x \in \mathbb{R}^m$. Identity (6.1) yields a representation of $(\sum_{i=1}^m x_i^2)^{q/2}$ as a sum of q th powers of real linear forms with positive real coefficients. Such a representation is called a *Hilbert identity* [25]. Various aesthetic meanings of Hilbert identities are extensively discussed in a famous paper by Reznick [27].

Many Hilbert identities can be obtained by the cubature that are constructed in Sections 4 and 5. In particular, some of the resulting identities are represented as a sum of q th powers of rational linear forms with positive rational coefficients. Such *rational representations* were used not only in studying Waring's problem [6, pp. 717-725], but also in the work of Schmid on real holomorphy rings [30]. An aesthetic meaning of rational representations would be stated as follows⁵: We would take all coefficients $\{a_i\}$ which appear in a formula, and consider the field created by adjoining them, and then look at its dimension $[\mathbb{Q}(\{a_i\}) : \mathbb{Q}]$. With this measure, the “best formulas” would only involve rationals, and the minimum value occurs if the coefficients are already in $\mathbb{Q}$.

It is well known (it goes back to Hilbert [16]) that

$$(6.2) \quad \int_{S^{m-1}} y_1^q \rho(dy) = \frac{(q-1)!!(m-2)!!}{(m+q-2)!!}.$$

This is certainly a rational number. All cubature given in Section 4 have rational weights, and points from orbits of the form $(\sqrt[q]{a}, \dots, \sqrt[q]{a}, 0, \dots, 0)^{B_m}$ with rational a . Thus, by Proposition 2.3, we can obtain many rational representations.

For example, the 91-point cubature of Example 4.9 is translated into the following rational representation that Reznick [27] was not able to find.

Theorem 6.1.

$$(6.3) \quad 120 \left(\sum_{i=1}^7 x_i^2 \right)^3 = \sum_{56} (x_i \pm x_{i+2} \pm x_{i+3} \pm x_{i+4})^6 + 2 \sum_{28} (x_i \pm x_{i+2} \pm x_{i+3})^6 + \sum_7 (2x_i)^6$$

where on the right the indices are taken as cyclic modulo 7 and all possible combinations of signs occur in the summation.

Remark 6.2. Reznick [27, p. 112] translated an index-six cubature on S^6 which was found by Stroud in 1967 into the following beautiful representation:

$$(6.4) \quad 960 \left(\sum_{i=1}^7 x_i^2 \right)^3 = 2 \sum_7 (2x_i)^6 + \sum_{2 \cdot \binom{7}{2}} (2x_i \pm 2x_j)^6 + \sum_{2^6} (x_1 \pm \dots \pm x_7)^6,$$

where on the right all possible combinations of signs and pairs of the 7 variables $x_1, \dots, x_7$ occur in the second summation. Identity (6.3) improves Reznick's representation. Namely, (6.3) has fewer number of sixth powers than (6.4).

⁵This was suggested by Bruce Reznick through email conversation.

More rational representations are available. For example, look at the following Kürschák's representation:

$$2^k \binom{3k}{k} \left(\sum_{i=1}^{3k+1} x_i^2 \right)^2 = \sum (x_{i_1} \pm x_{i_2} \pm \dots \pm x_{i_{k+1}})^4$$

where on the right all possible combinations of signs and $(k+1)$ -subsets of the $3k+1$ variables $x_1, \dots, x_{3k+1}$ occur [6, p. 723]. This corresponds to the cubature of Lemma 4.4 (ii), which is, by Theorem 4.3, reduced to many rational representations involving much fewer number of fourth powers.

We give one more interesting Hilbert identity, though it is not always rational.

Theorem 6.3.

$$\begin{aligned} \left(\sum_{i=1}^4 x_i^2 \right)^5 &= \frac{1}{2520} \sum_4 (2x_i)^{10} + \frac{1}{2520} \sum_8 (x_1 \pm x_2 \pm x_3 \pm x_4)^{10} \\ &\quad + \frac{1-120a}{272160} \sum_{32} (3x_i \pm x_j \pm x_k \pm x_l)^{10} + \frac{1-120a}{272160} \sum_{16} (2x_i \pm 2x_j \pm 2x_k)^{10} \\ (6.5) \quad &\quad + \frac{192a-1}{68040} \sum_{48} (2x_i \pm x_j \pm x_k)^{10} + \frac{12-960a}{630} \sum_{12} (x_i \pm x_j)^{10}, \end{aligned}$$

where $\frac{1}{192} \leq a \leq \frac{1}{120}$. In particular, if a is rational, then so is the corresponding identity.

Proof. The cubature of Theorem 5.10 (1) is centrally symmetric, which is reduced to the half-size formula of index 10. The result then follows by (6.1) and (6.2). $\square$

Identity (6.5) unifies the following familiar identity by I. Schur (cf. [6, p. 721]).

Corollary 6.4.

$$\begin{aligned} 22680 \left(\sum_{i=1}^4 x_i^2 \right)^5 &= 9 \sum_4 (2x_i)^{10} + 9 \sum_8 (x_1 \pm x_2 \pm x_3 \pm x_4)^{10} \\ (6.6) \quad &\quad + \sum_{48} (2x_i \pm x_j \pm x_k)^{10} + 180 \sum_{12} (x_i \pm x_j)^{10}. \end{aligned}$$

Proof. Take $a = 1/120$ in (6.5). $\square$

Remark 6.5. Some classical identities as such by Lucas (1876) and Liouville (1859), are often picked up for an introduction in the study of Hilbert identities [6]. It is well known (see, e.g., [14, 27]) that Liouville's and Lucas's identities are closely related by a linear change and provide essentially the same cubature on S^3 . The Hurwitz identity

$$\begin{aligned} 5040 \left(\sum_{i=1}^4 x_i^2 \right)^4 &= 6 \sum_4 (2x_i)^8 + 6 \sum_8 (x_1 \pm x_2 \pm x_3 \pm x_4)^8 \\ &\quad + \sum_{48} (2x_i \pm x_j \pm x_k)^8 + 60 \sum_{12} (x_i \pm x_j)^8 \end{aligned}$$

is also well known [6, p. 721]. It is interesting to note that Hurwitz's and Schur's identities are the same in terms of spherical cubature, i.e., the corresponding formulae have the same weights and points. In [14, 27], this observation is not remarked, though the relation between Liouville's and Lucas's identities is mentioned.

The story so far implies how powerful the cubature approach is to construct Hilbert identities. In turn, we look at an advantage of translating spherical cubature into Hilbert identities.

Theorem 6.6. *Let $m \geq 2$ be an integer. Then $(\sum_{i=1}^m x_i^2)^4$ does not have a representation as an $\mathbb{R}$ -linear combination of $(a_1x_1 + \cdots + a_mx_m)^8$ with $a_i \in \{0, -1, 1\}$.*

Proof. The ratio of the coefficients of $x_1^6x_2^2$ and $x_1^4x_2^4$ is $(2:3)$ in $(\sum_{i=1}^m x_i^2)^4$. But it is $(2:5)$ in any form $(a_1x_1 + \cdots + a_nx_n)^8$ with $a_i \in \{0, \pm 1\}$, $0 \notin \{a_1, a_2\}$. $\square$

Corollary 6.7. *Let $m \geq 2$ and G be a subgroup of B_m . Then there exists no G -invariant Euclidean 8-design of $\mathbb{R}^m$ that consists of the orbits of the form $(1, \dots, 1, 0, \dots, 0)^G$.*

Proof. Restricting (2.3) to homogeneous polynomials of degree 8 implies the existence of a cubature of index 8 on S^{m-1} , by suitably rescaling points and weights. The result then follows by Theorem 6.6. $\square$

A variation of Corollary 6.7 holds for all irreducible reflection groups. Namely, Theorem 5.1 can be proved even if each irreducible reflection group is replaced by its subgroups.

Remark 6.8. (i) Corollary 6.7 is the Bajnok theorem for $G = B_m$, and the case (3) of Theorem 5.1 for $G = D_m$. It is also interesting to note that Theorem 6.6 states that the Bajnok theorem is valid even if negative coefficients are allowed. (ii) To prove Theorem 2.10, Bajnok implicitly used the Sobolev theorem. The approach based on the Sobolev theorem is of theoretic interest, but it basically requires tedious calculations on invariant harmonic homogeneous polynomials. In summary, the original proof of Bajnok requires a few pages [1, Section 2 and Proposition 15] and seems to be involved. Whereas, the present proof is short, and simple for it uses only elementary counting techniques. The Bajnok theorem is well known in algebra and combinatorics, however, is not fully recognized in numerical analysis, though it can be used to determine the maximum degree of a symmetric cubature on the simplex [38] which is traditionally studied in the context of numerical analysis ⁶. The authors expect that the new proof will make researchers in many fields more familiar with the Bajnok theorem.

Acknowledgement This work started when the second author stayed at the Department of Mathematics of the University of Oregon from April to June in 2011. He gratefully acknowledges the hospitality of this institution and the cooperation with Yuan Xu and many other staffs. The authors also thank Eiichi Bannai, Reinhard Laue, Sanpei Kageyama, and Oksana Shatalov for fruitful discussions about regular t -wise balanced designs and index-type cubature. After an earlier version of this paper was written, the second author emailed Bruce Reznick and Koichi Kawada to discuss the contents of Sections 5 and 6. They were really patient in giving us some elementary courses in the subject and many valuable comments and suggestions, and the resulting revision extensively improved the previous version.

⁶ The second author learned this fact from Yuan Xu. In [29], we proved a variation of the Bajnok theorem for cubature formulae on the simplex, particularly intended for researchers in numerical analysis.

APPENDIX A. CLASSIFICATION OF E_8 -INVARIANT CUBATURE

An E_8 -invariant cubature of degree 15 that consists of the orbits of the corner vectors is classified by the following 27 types of weights:

$$\begin{aligned} w_1 &= \frac{23}{504000} - \frac{4288512w_2}{823543} - \frac{258048w_3}{15625} - \frac{70224w_4}{15625} - \frac{15w_8}{128}, \\ w_5 &= \frac{3}{224000} - \frac{1244160w_2}{823543} - \frac{171008w_3}{15625} - \frac{79704w_4}{15625} - \frac{243w_8}{512}, \\ w_6 &= \frac{9}{896000} + \frac{4193208w_2}{823543} + \frac{507384w_3}{15625} + \frac{180792w_4}{15625} + \frac{3645w_8}{2048}, \\ w_7 &= \frac{67}{672000} - \frac{2465280w_2}{823543} - \frac{290304w_3}{15625} - \frac{94752w_4}{15625} - \frac{603w_8}{512}, \end{aligned}$$

and

- (1) $w_4 = 0, 0 \leq w_2 \leq \frac{12588443}{1449551462400}, 0 \leq w_3 < \frac{44118375-4976640000000w_2}{36053104984064},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3}{3126889828125},$
- (2) $w_4 = 0, 0 \leq w_2 \leq \frac{12588443}{1449551462400}, w_3 = \frac{44118375-4976640000000w_2}{36053104984064}, w_8 = 0,$
- (3) $w_4 = 0, \frac{12588443}{1449551462400} < w_2 < \frac{117649}{13436928000}, 0 \leq w_3 \leq \frac{44118375-5038848000000w_2}{14396954722304},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3}{3126889828125},$
- (4) $w_4 = 0, \frac{12588443}{1449551462400} < w_2 < \frac{117649}{13436928000},$
 $\frac{44118375-5038848000000w_2}{14396954722304} < w_3 < \frac{338240875-38596608000000w_2}{122407847460864},$
 $0 \leq w_8 \leq \frac{676481750-77193216000000w_2-244815694921728w_3}{1737161015625},$
- (5) $w_4 = 0, \frac{12588443}{1449551462400} < w_2 < \frac{117649}{13436928000},$
 $w_3 = \frac{338240875-38596608000000w_2}{122407847460864}, w_8 = 0,$
- (6) $w_4 = 0, w_2 = \frac{117649}{13436928000}, w_3 = 0, 0 \leq w_8 \leq \frac{2}{5740875},$
- (7) $w_4 = 0, w_2 = \frac{117649}{13436928000}, 0 < w_3 < \frac{625}{252829237248}, 0 \leq w_8 \leq \frac{1250-505658474496w_3}{3588046875},$
- (8) $w_4 = 0, w_2 = \frac{117649}{13436928000}, w_3 = \frac{625}{252829237248}, w_8 = 0,$
- (9) $w_4 = 0, \frac{117649}{13436928000} < w_2 < \frac{2705927}{308772864000}, 0 \leq w_3 < \frac{338240875-38596608000000w_2}{122407847460864},$
 $0 \leq w_8 \leq \frac{676481750-77193216000000w_2-244815694921728w_3}{1737161015625},$
- (10) $w_4 = 0, \frac{117649}{13436928000} < w_2 < \frac{2705927}{308772864000}, w_3 = \frac{338240875-38596608000000w_2}{122407847460864},$
 $w_8 = 0,$
- (11) $w_4 = 0, w_2 = \frac{2705927}{308772864000}, w_3 = 0, w_8 = 0,$
- (12) $0 < w_4 < \frac{125}{3092173056}, 0 \leq w_2 < \frac{62942215+7929414597888w_4}{7247757312000},$
 $0 \leq w_3 < \frac{44118375-4976640000000w_2-16803755845632w_4}{36053104984064},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3-33607511691264w_4}{3126889828125},$
- (13) $0 < w_4 < \frac{125}{3092173056}, 0 \leq w_2 < \frac{62942215+7929414597888w_4}{7247757312000},$
 $w_3 = \frac{44118375-4976640000000w_2-16803755845632w_4}{36053104984064}, w_8 = 0,$
- (14) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{62942215+7929414597888w_4}{7247757312000},$
 $0 \leq w_3 < \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3-33607511691264w_4}{3126889828125},$
- (15) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{62942215+7929414597888w_4}{7247757312000},$
 $w_3 = \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864}, w_8 = 0,$
- (16) $0 < w_4 < \frac{125}{3092173056}, \frac{62942215+7929414597888w_4}{7247757312000} < w_2 < \frac{14706125-1123879249584w_4}{1679616000000},$
 $0 \leq w_3 \leq \frac{44118375-5038848000000w_2-3371637748752w_4}{14396954722304},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3-33607511691264w_4}{3126889828125},$
- (17) $0 < w_4 < \frac{125}{3092173056}, \frac{62942215+7929414597888w_4}{7247757312000} < w_2 < \frac{14706125-1123879249584w_4}{1679616000000},$
 $\frac{375}{122372096} - \frac{307546875w_2}{878720381} - \frac{255879w_4}{1092608} < w_3 < \frac{2875}{1040449536} - \frac{5453125w_2}{17294403} - \frac{209w_4}{768},$
 $0 \leq w_8 \leq \frac{676481750-77193216000000w_2-244815694921728w_3-66623021144064w_4}{1737161015625},$

- (18) $0 < w_4 < \frac{125}{3092173056}, \frac{62942215+7929414597888w_4}{7247757312000} < w_2 < \frac{14706125-1123879249584w_4}{1679616000000},$
 $w_3 = \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864}, w_8 = 0,$
- (19) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{14706125-1123879249584w_4}{1679616000000}, w_3 = 0,$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-33607511691264w_4}{3126889828125},$
- (20) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{14706125-1123879249584w_4}{1679616000000},$
 $0 < w_3 < \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864},$
 $0 \leq w_8 \leq \frac{676481750-77193216000000w_2-244815694921728w_3-66623021144064w_4}{1737161015625},$
- (21) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{14706125-1123879249584w_4}{1679616000000},$
 $w_3 = \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864}, w_8 = 0,$
- (22) $0 < w_4 < \frac{125}{3092173056}, \frac{14706125-1123879249584w_4}{1679616000000} < w_2 < \frac{338240875-33311510572032w_4}{38596608000000},$
 $0 \leq w_3 < \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864},$
 $0 \leq w_8 \leq \frac{676481750-77193216000000w_2-244815694921728w_3-66623021144064w_4}{1737161015625},$
- (23) $0 < w_4 < \frac{125}{3092173056}, \frac{14706125-1123879249584w_4}{1679616000000} < w_2 < \frac{338240875-33311510572032w_4}{38596608000000},$
 $w_3 = \frac{338240875-38596608000000w_2-33311510572032w_4}{122407847460864}, w_8 = 0,$
- (24) $0 < w_4 < \frac{125}{3092173056}, w_2 = \frac{338240875-33311510572032w_4}{38596608000000}, w_3 = 0, w_8 = 0,$
- (25) $\frac{125}{3092173056} \leq w_4 \leq \frac{125}{47609856}, 0 \leq w_2 < \frac{14706125-5601251948544w_4}{1658880000000},$
 $0 \leq w_3 < \frac{44118375-4976640000000w_2-16803755845632w_4}{36053104984064},$
 $0 \leq w_8 \leq \frac{88236750-9953280000000w_2-72106209968128w_3-33607511691264w_4}{3126889828125},$
- (26) $\frac{125}{3092173056} \leq w_4 \leq \frac{125}{47609856}, 0 \leq w_2 < \frac{14706125-5601251948544w_4}{1658880000000},$
 $w_3 = \frac{44118375-4976640000000w_2-16803755845632w_4}{36053104984064}, w_8 = 0,$
- (27) $\frac{125}{3092173056} \leq w_4 \leq \frac{125}{47609856}, w_2 = \frac{14706125-5601251948544w_4}{1658880000000}, w_3 = 0, w_8 = 0.$

REFERENCES

- [1] B. BAJNOK. *Orbits of the hyperoctahedral group as Euclidean designs*. J. Algebr. Comb. **25** (2007), 375–397.
- [2] EI. BANNAL, ETSU. BANNAL. *Tight Gaussian 4-designs*. J. Algebr. Comb. **22** (2005), 39–63.
- [3] EI. BANNAL, R. M. DAMERELL. *Tight spherical designs II*. J. London Math. Soc. **21** (1980), 13–30.
- [4] N. BOURBAKI. *Lie Groups and Lie Algebras: Chapters 4-6* (Elements of Mathematics). Springer, 2002.
- [5] P. DELSARTE, J. M. GOETHALS, J. J. SEIDEL. *Spherical codes and designs*. Geom. Dedicata **6** (1977), 363–388.
- [6] L. E. DICKSON. *History of the Theory of Numbers, II*. Carnegie Institution of Washington, 1923.
- [7] W. J. ELLISON. *Waring’s problem*. Amer. Math. Monthly **78** (1971), 10–35.
- [8] R. FUJI-HARA, S. KURIKI, M. JIMBO. *On balanced complementation for regular t -wise balanced designs*. Discrete Math. **76** (1989), 29–35.
- [9] J. M. GOETHALS, J. J. SEIDEL. *Cubature formulae, polytopes, and spherical designs*, in: Geometric Vein, The Coxeter Festschrift, Springer, Berlin, 1981, pp. 203–218.
- [10] S. C. GUPTA, B. JONES. *Equireplicate balanced block designs and unequal block sizes*. Biometrika **70** (1983), 433–440.
- [11] R. H. HARDIN, N. J. A. SLOANE. *Expressing $(a^2 + b^2 + c^2 + d^2)^3$ as a sum of 23 sixth powers*. J. Combin. Theory Ser. A **68** (1994), 481–485.
- [12] R. H. HARDIN, N. J. A. SLOANE. *McLaren’s improved snub cube and other new spherical designs in three dimensions*. Discrete Comput. Geom. **15** (1996), 429–441.
- [13] P. DE LA HARPE, C. PACHE, B. VENKOV. *Construction of spherical cubature formulas using lattices*. St. Petersburg Math. J. **18** (2007), 119–139.
- [14] P. DE LA HARPE, C. PACHE. *Cubature Formulas, Geometrical Designs, Reproducing Kernels, and Markov Operators*. Infinite Groups: Geometric, Combinatorial and Dynamical Aspects Progress in Mathematics, 2005, Vol. 248, 219–267.

- [15] A. S. HEDAYAT, N. J. A. SLOANE, J. STUFKEN. *Orthogonal Arrays. Theory and Applications*. Springer Series in Statistics, 1999.
- [16] D. HILBERT. *Beweis für die Darstellbarkeit der ganzen Zahlen durch eine feste Anzahl n -ter Potenzen (Waringsches Problem)*. Math. Ann. **67** (1909), 281–300.
- [17] Y. J. IONIN, T. TRUNG. *Symmetric Designs*, in: C. J. Colbourn, J. H. Dinitz (eds.), Handbook of Combinatorial Designs (2nd ed.). CRC Press, Boca Raton, USA, 2007, pp. 110–124.
- [18] S. KAGEYAMA, D. MAJUMDAR. *Resistant BTIB designs*. Commun. Statist. Theory Methods **19** (1990), 2145–2158.
- [19] G. B. KHOSROVSHAHI, R. LAUE. *t -Designs with $t \geq 3$* , in: C. J. Colbourn, J. H. Dinitz (eds.), Handbook of Combinatorial Designs (2nd ed.). CRC Press, Boca Raton, USA, 2007, pp. 79–101.
- [20] H. KÖNIG. *Isometric embeddings of Euclidean spaces into finite-dimensional l_p -spaces*, in: Banach Center Publ. **34**, Polish Acad. Sci. Warsaw, 1995, pp. 79–87.
- [21] T. LYONS, N. VICTOIR. *Cubature on Wiener space. Stochastic analysis with applications to mathematical finance*. Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **460** (2004), 169–198.
- [22] Y. I. LYUBICH, L. N. VASERSTEIN. *Isometric embeddings between classical Banach spaces, cubature formulas, and spherical designs*. Geom. Dedicata **47** (1993), 327–362.
- [23] A. NEUMAIER, J. J. SEIDEL. *Discrete measures for spherical designs, eutactic stars and lattices*. Nederl. Akad. Wetensch. Proc. Ser. A 91=Indag. Math. **50** (1988), 321–334.
- [24] H. NOZAKI, M. SAWA. *Note on cubature formulae and designs obtained from group orbits*. To appear in Canad. J. Math. (arXiv:1102.4891v1).
- [25] B. REZNICK. *On the length of binary forms*, in: K. Alladi, M. Bhargava, D. Saritt, P. Tiep (eds.), Developments in Math., Springer, New York (<http://arxiv.org/pdf/1007.5485.pdf>).
- [26] B. REZNICK. *Some constructions of spherical 5-designs*. Linear Algebra Appl. **226-228** (1995), 163–196.
- [27] B. REZNICK. *Sums of even powers of real linear forms*. Mem. Amer. Math. Soc. **96** (1992), No. 463.
- [28] G. N. SALIHOV. *Cubature formulas for a hypersphere that are invariant with respect to the group of the regular 600-face*. Soviet Math. Dokl. **16** (1975), 1046–1050.
- [29] M. SAWA, Y. XU. *On positive cubature rules on the simplex and isometric embeddings*. arXiv:1108.3385v1.
- [30] J. SCHMID. *On totally positive units of real holomorphy rings*. Israel J. Math. **85** (1994), 339–350.
- [31] J. J. SEIDEL. *Isometric embeddings and geometric designs*. Discrete Math. **136** (1994), 281–293.
- [32] O. SHATALOV. *Isometric embeddings $l_2^m \rightarrow l_p^n$ and cubature formulas over classical fields*. Doctor Thesis, Technion-Israel Institute of Technology, Haifa, Israel, 2001.
- [33] S. L. SOBOLEV. *Cubature formulas on the sphere which are invariant under transformations of finite rotation groups* (in Russian). Dokl. Akad. Nauk SSSR **146** (1962), 310–313.
- [34] A. H. STROUD. *Approximate Calculation of Multiple Integrals*. Prentice-Hall, Englewood Cliffs, N.J., 1971.
- [35] N. VICTOIR. *Asymmetric cubature formulae with few points in high dimension for symmetric measures*. SIAM J. Numer. Anal. **42** (2004), 209–227.
- [36] W. G. WOODAL. *Square λ -linked designs*. Proc. London Math. Soc. **20** (1970), 669–687.
- [37] Y. XU. *Minimal cubature formulae for a family of radial weight functions*. Adv. Comput. Math. **8** (1998), 367–380.
- [38] Y. XU. *Orthogonal polynomials and cubature formulae on spheres and on simplices*. Methods Appl. Anal. **5** (1998), 169–184.

DEPARTMENT OF MATHEMATICS, AICHI UNIVERSITY OF EDUCATION, IGAYA-CHO, KARIYA-CITY 448-8542, JAPAN

E-mail address: hnozaki@aeu.ac.jp

GRADUATE SCHOOL OF INFORMATION SCIENCES, NAGOYA UNIVERSITY, CHIKUSA-KU, NAGOYA 464-8601, JAPAN

E-mail address: sawa@is.nagoya-u.ac.jp